

# MINIMALITY AND MUTATION-EQUIVALENCE OF POLYGONS

ALEXANDER M. KASPRZYK, THOMAS PRINCE, AND BENJAMIN NILL

ABSTRACT. We introduce a concept of minimality for Fano polygons. We show that, up to mutation, there are only finitely many Fano polygons with given singularity content, and give an algorithm to determine representatives for all mutation-equivalence classes of such polygons. This is a key step in a program to classify orbifold del Pezzo surfaces using mirror symmetry. As an application, we classify all Fano polygons such that the corresponding toric surface is qG-deformation-equivalent to either (i) a smooth surface; or (ii) a surface with only singularities of type  $\frac{1}{3}(1, 1)$ .

## 1. INTRODUCTION

**1.1. An introduction from the viewpoint of algebraic geometry and mirror symmetry.** A *Fano polygon*  $P$  is a convex polytope in  $N_{\mathbb{Q}} := N \otimes_{\mathbb{Z}} \mathbb{Q}$ , where  $N$  is a rank-two lattice, with primitive vertices  $\mathcal{V}(P)$  in  $N$  such that the origin is contained in its strict interior,  $\mathbf{0} \in P^{\circ}$ . A Fano polygon defines a toric surface  $X_P$  given by the *spanning fan* (also commonly referred to as the *face fan* or *central fan*) of  $P$ ; that is,  $X_P$  is defined by the fan whose cones are spanned by the faces of  $P$ . The toric surface  $X_P$  has cyclic quotient singularities (corresponding to the cones over the edges of  $P$ ) and the anti-canonical divisor  $-K_{X_P}$  is  $\mathbb{Q}$ -Cartier and ample. Hence  $X_P$  is a toric *del Pezzo surface*.

The simplest example of a toric del Pezzo surface is  $\mathbb{P}^2$ , corresponding, up to  $GL_2(\mathbb{Z})$ -equivalence, to the triangle  $P = \text{conv}\{(1, 0), (0, 1), (-1, -1)\}$ . It is well-known that there are exactly five smooth toric del Pezzo surfaces, and that these are a subset of the sixteen toric Gorenstein del Pezzo surfaces (in bijective correspondence with the famous sixteen reflexive polygons described by Batyrev and Rabinowitz [7, 37]). More generally, if one bounds the *Gorenstein index*  $r$  (the smallest positive integer such that  $-rK_{X_P}$  is Cartier) the number of possibilities is finite. Dais classified those toric del Pezzo surfaces with Picard rank one and  $r \leq 3$  [15]. A general classification algorithm was presented in [28].

A new viewpoint on del Pezzo classification is suggested by mirror symmetry. We shall sketch this briefly; for details see [11]. An  $n$ -dimensional Fano variety  $X$  is expected to correspond, under mirror symmetry, to a Laurent polynomial  $f \in \mathbb{C}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$  [5, 8, 11, 12]. Under this correspondence, the regularised quantum period  $\widehat{G}_X$  of  $X$  – a generating function for Gromov–Witten invariants – coincides with the classical period  $\pi_f$  of  $f$  – a solution of the associated Picard–Fuchs differential equation – given by

$$\pi_f(t) = \left( \frac{1}{2\pi i} \right)^n \int_{|x_1|=\dots=|x_n|=1} \frac{1}{1 - tf(x_1, \dots, x_n)} \frac{dx_1}{x_1} \cdots \frac{dx_n}{x_n} = \sum_{k \geq 0} \text{coeff}_1(f^k) t^k.$$

If a Fano variety  $X$  is mirror to a Laurent polynomial  $f$  then it is expected that  $X$  admits a degeneration to the singular toric variety  $X_P$  associated to the Newton polytope  $P$  of  $f$ .

In general there will be many (often infinitely many) different Laurent polynomials mirror dual to  $X$ , and hence many toric degenerations  $X_P$ . It is conjectured that these Laurent polynomials are related via birational transformations analogous to cluster transformations, which are called *mutations* [2, 17, 19, 20]. A mutation acts on the Newton polytope  $P := \text{Newt}(f) \subset N_{\mathbb{Q}}$  of a Laurent polynomial via “rearrangement of Minkowski slices” (see §2.1), and on the dual polytope  $P^* \subset M_{\mathbb{Q}}$ ,  $M := \text{Hom}(N, \mathbb{Z})$ , via a piecewise- $GL_n(\mathbb{Z})$  transformation (see §2.2) [2]. At the level of Laurent polynomials, if  $f$  and  $g$  are related via mutation then their classical periods agree [2, Lemma 2.8]:  $\pi_f = \pi_g$ . Ilten [26] has shown that if two Fano polytopes  $P$  and  $Q$  are related by mutation then the

2010 *Mathematics Subject Classification*. 14J45 (Primary); 14J10, 14M25, 52B20 (Secondary).

*Key words and phrases*. Del Pezzo surface, lattice polygon, Fano polygon, mutation, cluster algebra.

corresponding toric varieties  $X_P$  and  $X_Q$  are deformation equivalent: there exists a flat family  $\mathcal{X} \rightarrow \mathbb{P}^1$  such that  $\mathcal{X}_0 \cong X_P$  and  $\mathcal{X}_\infty \cong X_Q$ . In fact  $X_P$  and  $X_Q$  are related via a  $\mathbb{Q}$ -Gorenstein (qG) deformation [1].

Classifying Fano polygons up to mutation-equivalence thus becomes a fundamental problem. One important mutation invariant is the *singularity content* (see §3.1) [3]. This consists of a pair  $(n, \mathcal{B})$ , where  $n$  is an integer – the number of *primitive T-singularities* – and  $\mathcal{B}$  is a *basket* – a collection of so-called *residual singularities*. A residual singularity is a cyclic quotient singularity that is rigid under qG-deformations; at the other extreme a *T-singularity* is a cyclic quotient singularity that admits a qG-smoothing [29]. The toric del Pezzo surface  $X_P$  is qG-deformation-equivalent to a del Pezzo surface  $X$  with singular points given by  $\mathcal{B}$  and Euler number of the non-singular locus  $X \setminus \text{Sing}(X)$  equal to  $n$ .

**Definition 1.1** ([1]). A del Pezzo surface with cyclic quotient singularities that admits a qG-degeneration (with reduced fibres) to a normal toric del Pezzo surface is said to be of class *TG*.

Notice that not all del Pezzo surfaces can be of class *TG*: not every del Pezzo has  $h^0(X, -K_X) > 0$ , for example (see Example 3.9). But it is natural to conjecture the following:

**Conjecture A** ([1]). There exists a bijective correspondence between the set of mutation-equivalence classes of Fano polygons and the set of qG-deformation-equivalence classes of locally qG-rigid class *TG* del Pezzo surfaces with cyclic quotient singularities.

The main results of this paper can be seen as strong evidence in support of the conjecture above. First, an immediate consequence of Theorem 5.6 is:

**Theorem 1.2.** *There are precisely ten mutation-equivalence classes of Fano polygons such that the toric del Pezzo surface  $X_P$  has only T-singularities. They are in bijective correspondence with the ten families of smooth del Pezzo surfaces.*

Second, combining the results of [14] with Theorem 7.4 we have:

**Theorem 1.3.** *There are precisely 26 mutation-equivalence classes of Fano polygons with singularity content  $(n, \{m \times \frac{1}{3}(1, 1)\})$ ,  $m \geq 1$ . They are in bijective correspondence with the 26 qG-deformation families of del Pezzo surfaces with  $m \times \frac{1}{3}(1, 1)$  singular points that admit a toric degeneration.*

In Theorem 6.3 we prove more generally that, up to mutation, the number of Fano polygons with basket  $\mathcal{B}$  is finite, and give an algorithm that outputs a finite list of Fano polygons that contains representatives for all their mutation-equivalence classes. It may happen that some of the Fano polygons produced by the algorithm are representatives for the same mutation-equivalence class. While there is currently no algorithm to determine whether two Fano polygons are mutation-equivalent, a useful necessary condition is provided by Lemma 3.11. If one accepts Conjecture A then Theorem 6.3 tells us that, for fixed basket  $\mathcal{B}$ , the number of qG-deformation-equivalence classes of del Pezzo surfaces of type *TG* with singular points  $\mathcal{B}$  is finite, and gives an algorithm for classifying their toric degenerations.

**1.2. An introduction from the viewpoint of cluster algebras and cluster varieties.** One can obtain information about mutations of polygons using quivers and the theory of cluster algebras [16, 17]. There is a precise analogy between mutation classes of Fano polygons and the clusters of certain cluster algebras, as we now describe. Let  $L \cong \mathbb{Z}^n$ , and fix a skew-symmetric form  $\{\cdot, \cdot\}$  on  $L$ . A *cluster*  $C$  is a transcendence basis for  $\mathbb{C}(L)$ , and a *seed* is a pair  $(B, C)$  where  $B$  is a basis of  $L$ . There is a notion of *mutation* of seeds, given in Definition 3.14 below; this depends on the form  $\{\cdot, \cdot\}$ . A *cluster algebra* is the algebra generated by all clusters that can be obtained from a given initial seed by mutation. To a seed  $(B, C)$  one can associate a quiver  $Q_B$  with vertex set equal to  $B$  and the number of arrows from  $e_i \in B$  to  $e_j \in B$  equal to  $\max(\{e_i, e_j\}, 0)$ . Changing the seed  $(B, C)$  by a mutation changes the quiver  $Q_B$  by a *quiver mutation* (see Definition 3.16). Conversely, from a quiver with vertex set  $B$  and no vertex-loops or two-cycles, one can construct a cluster algebra by setting  $L = \mathbb{Z}^B$ , defining  $\{e_i, e_j\}$  to be the (signed) number of arrows from  $e_i \in B$  to  $e_j \in B$ , and taking the initial seed to be  $(B, C)$ , where  $C$  is the standard transcendence basis for  $\mathbb{C}(L)$ .

We can also associate a quiver and a cluster algebra to a Fano polygon  $P$  as follows. Suppose that the singularity content of  $P$  is  $(n, \mathcal{B})$ . The associated quiver  $Q_P$  has  $n$  vertices; each vertex  $v$  corresponds

to a *primitive  $T$ -singularity* (Definition 3.1), and hence determines an edge  $E$  of  $P$  (these edges need not be distinct). The number of arrows from vertex  $v$  to vertex  $v'$  is defined to be  $\max(w \wedge w', 0)$ , where  $\wedge$  denotes the determinant, and  $w$  and  $w'$  are the primitive inner normal vectors to the edges  $E$  and  $E'$  of  $P$ . The cluster algebra  $\mathcal{A}_P$  associated to  $P$  is the cluster algebra associated to  $Q_P$ ; we denote the initial seed of this cluster algebra by  $(B_P, C_P)$ .

We show in Proposition 3.15 below that a mutation from a seed  $(B, C)$  to a seed  $(B', C')$  induces a mutation between the corresponding Fano polygons  $P$  and  $P'$ . We then show, in Proposition 3.17, that a mutation from a Fano polygon  $P$  to a Fano polygon  $P'$  induces a mutation between the corresponding quivers  $Q_P$  and  $Q_{P'}$ . These correspondences have consequences for mutation-equivalence which are not readily apparent from the polygon alone.

**Example 1.4.** Consider a Fano polygon  $P \subset N_{\mathbb{Q}}$  containing only two primitive  $T$ -singularities, and suppose that the corresponding inner normal vectors form a basis for the dual lattice  $M$ . Then there are at most five polygons, up to  $GL_2(\mathbb{Z})$ -equivalence, that are mutation-equivalent to  $P$ . This follows from the fact that the quiver associated to  $P$  has underlying graph  $A_2$ , and that the exchange graph of the  $A_2$  cluster algebra is pentagonal; see Corollary 3.18 below.

**1.3. An introduction from the viewpoint of the geometry of numbers.** The relation between the lattice points in a convex body and its geometric shape and volume is a key problem in convex geometry and integer optimisation. These connections have been addressed specifically for lattice polytopes, independently of their significance in toric geometry. Here we focus only on the case of interest in this paper, that of a Fano polygon. A classical result in this area is the following (these statements can be generalised and quantified, see [24]):

**Theorem 1.5** ([31, 39]). *There are only finitely many  $GL_2(\mathbb{Z})$ -equivalence classes of Fano polygons  $P$  with  $I$  interior lattice points, for each  $I \in \mathbb{Z}_{>0}$ .*

**Corollary 1.6.** *There are only a finite number of possibilities for the area and number of lattice points of a Fano polygon with  $I$  interior lattice points, for each  $I \in \mathbb{Z}_{>0}$ .*

In [2] a new equivalence relation on Fano polygons  $P$  was introduced, called *mutation-equivalence*, that is weaker than  $GL_2(\mathbb{Z})$ -equivalence. In particular there exist infinitely many mutation-equivalent Fano polygons that are not  $GL_2(\mathbb{Z})$ -equivalent (see, for example, [4, Example 3.14]) and so their area and number of lattice points cannot be bounded. Mutation-equivalence does, however, preserve the Ehrhart series (and hence area) of the dual polygon  $P^* \subset M_{\mathbb{Q}}$  (see §2.2) [2, Proposition 4].

**Remark 1.7.** It is known that the product  $\text{Vol}(P) \cdot \text{Vol}(P^*)$  of the (lattice normalised) area of a polygon  $P$  and of its dual polygon  $P^*$  cannot be arbitrarily small [32], however it is easy to construct families of polygons where both  $\text{Vol}(P)$  and  $\text{Vol}(P^*)$  can simultaneously become arbitrarily large. For example,  $P_k := \text{conv}\{(k, 1), (k, -1), (-1, 0)\}$  for  $k \in \mathbb{Z}_{>0}$ .

Given a Fano polygon  $P$  there exists an explicit formula for the Ehrhart series and area of the dual polygon  $P^*$  in terms of the *singularity content* of  $P$  (see §3.1) [3]. A consequence of this formula is that mutation-equivalent Fano polygons cannot have an arbitrarily large number of vertices [3, Lemma 3.8]; equivalently, singularity content gives an upper bound on the Picard rank of  $X_P$  under qG-deformation.

Recall that the *height*  $r_E \in \mathbb{Z}_{\geq 0}$  of a lattice line segment  $E \subset N_{\mathbb{Q}}$  is the lattice distance of  $E$  from the origin, and the *length* of the line segment  $E$  is given by the positive integer  $k = |E \cap N| - 1$ . Clearly there exist unique non-negative integers  $n$  and  $k_0$ ,  $0 \leq k_0 < r_E$ , such that  $k = nr_E + k_0$ . Suppose that  $E$  is the edge of a Fano polygon, so that the vertices of  $E$  are primitive. As described in [3] (see also §3.1), one can decompose  $E$  into  $n + 1$  (or  $n$  if  $k_0 = 0$ ) lattice line segments with primitive vertices. Of these,  $n$  line segments have their length equal to their height; the cones over these line segments correspond to *primitive  $T$ -singularities*. If  $k_0 \neq 0$  then there is one additional lattice line segment of length  $k_0 < r_E$ ; the cone over this line segment corresponds to a *residual singularity*.

Although there may be several different decompositions of this form for an edge  $E$ , it turns out that the residual singularity is unique – it does not depend on the choice of decomposition. In addition, the collection of residual singularities arising from all of the edges of  $P$ , which we call the *basket* of  $P$

and denote by  $\mathcal{B}$ , is a mutation invariant. We say that a lattice point of  $P$  is *residual* if it lies in the strict interior of a residual cone, for some fixed choice of decomposition. The number of residual lattice points does not depend on the chosen decomposition, but only on the basket  $\mathcal{B}$  of  $P$ , and is invariant under mutation. The main results of this paper can now be stated in a way analogous to the classical results above:

**Theorem 1.8.** *There are only finitely many mutation-equivalence classes of Fano polygons  $P$  with  $N$  residual lattice points, for each  $N \in \mathbb{Z}_{\geq 0}$ .*

*Proof.* If there are no residual cones then the result follows from Theorem 5.4. In order to use Theorem 6.3 we only have to show that the height  $r_E$  of an edge  $E$  containing a residual cone is bounded. Let  $v_1$  and  $v_2$  be primitive points on  $E$  such that  $\text{cone}\{v_1, v_2\}$  is a residual cone. The line segment joining  $v_1$  and  $v_2$  has length  $1 \leq k < r_E$ . We see that the lattice triangle

$$\text{conv}\{\mathbf{0}, v_1, v_1 + (v_2 - v_1)/k\}$$

has at most  $N + 3$  lattice points. Pick's formula [35] implies that its area and thus the height  $r_E$  is bounded in terms of  $N$ .  $\square$

**Corollary 1.9.** *There are only a finite number of possibilities for the dual area and the number of vertices of a Fano polygon with  $N$  residual lattice points, for each  $N \in \mathbb{Z}_{\geq 0}$ .*

In particular, this shows that there exist no Fano polygons with empty basket but an arbitrarily large number of vertices. Note that it can be easily seen that there exist centrally-symmetric Fano polygons with an arbitrarily large number of vertices where every edge corresponds to a residual singularity.

## 2. MUTATION OF FANO POLYGONS

In [2, §3] the concept of mutation for a lattice polytope was introduced. We state it here in the simplified case of a Fano polygon  $P \subset N_{\mathbb{Q}}$  and refer to [2] for the general definitions.

**2.1. Mutation in  $N$ .** Let  $w \in M := \text{Hom}(N, \mathbb{Z})$  be a primitive inner normal vector for an edge  $E$  of  $P$ , so  $w : N \rightarrow \mathbb{Z}$  induces a grading on  $N_{\mathbb{Q}}$  and  $w(v) = -r_E$  for all  $v \in E$ , where  $r_E$  is the height of  $E$ . Define

$$h_{\max} := \max\{w(v) \mid v \in P\} \quad \text{and} \quad h_{\min} := -r_E = \min\{w(v) \mid v \in P\}.$$

We have that  $h_{\max} > 0$  and  $h_{\min} < 0$ . For each  $h \in \mathbb{Z}$  we define  $w_h(P)$  to be the (possibly empty) convex hull of those lattice points in  $P$  at height  $h$ ,

$$w_h(P) := \text{conv}\{v \in P \cap N \mid w(v) = h\}.$$

By definition  $w_{h_{\min}}(P) = E$  and  $w_{h_{\max}}(P)$  is either a vertex or an edge of  $P$ . Let  $v_E \in N$  be a primitive lattice element of  $N$  such that  $w(v_E) = 0$ , and define  $F := \text{conv}\{\mathbf{0}, v_E\}$ , a line segment of unit length parallel to  $E$  at height 0. Notice that  $v_E$ , and hence  $F$ , is uniquely defined only up to sign.

**Definition 2.1.** Suppose that for each negative height  $h_{\min} \leq h < 0$  there exists a (possibly empty) lattice polytope  $G_h \subset N_{\mathbb{Q}}$  satisfying

$$(2.1) \quad \{v \in \mathcal{V}(P) \mid w(v) = h\} \subseteq G_h + |h|F \subseteq w_h(P),$$

where  $+$  denotes the Minkowski sum, and we define  $\emptyset + Q = \emptyset$  for any polytope  $Q$ . We call  $F$  a *factor* of  $P$  with respect to  $w$ , and define the *mutation* given by the primitive normal vector  $w$ , factor  $F$ , and polytopes  $\{G_h\}$  to be:

$$\text{mut}_w(P, F) := \text{conv} \left( \bigcup_{h=h_{\min}}^{-1} G_h \cup \bigcup_{h=0}^{h_{\max}} (w_h(P) + hF) \right) \subset N_{\mathbb{Q}}.$$

Although not immediately obvious from the definition, the resulting mutation is independent of the choices of  $\{G_h\}$  [2, Proposition 1]. Furthermore, up to isomorphism, mutation does not depend on the choice of  $v_E$ : we have that  $\text{mut}_w(P, F) \cong \text{mut}_w(P, -F)$ . Since we consider a polygon to be defined only up to  $GL_2(\mathbb{Z})$ -equivalence, mutation is well-defined and unique. Any mutation can be inverted

by inverting the sign of  $w$ : if  $Q := \text{mut}_w(P, F)$  then  $P = \text{mut}_{-w}(Q, F)$  [2, Lemma 2]. Finally, we note that  $P$  is a Fano polygon if and only if the mutation  $Q$  is a Fano polygon [2, Proposition 2].

We call two polygons  $P$  and  $Q \subset N_{\mathbb{Q}}$  *mutation-equivalent* if there exists a finite sequence of mutations between the two polygons (considered up to  $GL_2(\mathbb{Z})$ -equivalence). That is, if there exist polygons  $P_0, P_1, \dots, P_n$  with  $P \cong P_0$ ,  $P_{i+1} = \text{mut}_{w_i}(P_i, F_i)$ , and  $Q \cong P_n$ , for some  $n \in \mathbb{Z}_{\geq 0}$ .

**Remark 2.2.** We remark briefly upon the three ways in which our definition above differs slightly from that in [2].

- (i) First, [2] does not require that the factor  $F$  be based at the origin. The condition that  $\mathbf{0} \in \mathcal{V}(F)$  is harmless, and indeed we have touched on this above when we noted that  $F$  and  $-F$  give  $GL_2(\mathbb{Z})$ -equivalent mutations: in general translation of a factor  $F$  by a lattice point  $v \in w^{\perp} \cap N$ , where  $w^{\perp} := \{v \in N_{\mathbb{Q}} \mid w(v) = 0\}$ , results in isomorphic mutations. It is reasonable to regard a factor as being defined only up to translation by elements in  $w^{\perp} \cap N$ , with the resulting mutation defined only up to “shear transformations” fixing the points in  $w^{\perp}$ .
- (ii) Second, the more general definition places no restriction on the dimension of the factor  $F$ , although the requirement that  $F \subset w^{\perp}$  does mean that  $\text{codim}(F) \geq 1$ . In particular it is possible to take  $F = v$ , where  $w(v) = 0$ . But observe that  $v = \mathbf{0} + v$ , and  $\text{mut}_w(\cdot, \mathbf{0})$  is the identity, hence  $\text{mut}_w(P, v)$  is trivial. Thus our insistence that  $\dim(F) = 1$  is reasonable.
- (iii) Finally, our requirement that  $F$  is of unit length is a natural simplification: in general, if the factor can be written as a Minkowski sum  $F = F_1 + F_2$ , where we can insist that each  $F_i \subset w^{\perp}$  and  $\dim(F_i) > 0$ , then the mutation with factor  $F$  can be written as the composition of two mutations with factors  $F_1$  and  $F_2$  (with fixed  $w$ ). Thus it is reasonable to assume that the factor is Minkowski-indecomposable and hence, for us, a primitive line segment.

In two dimensions, mutations are completely determined by the edges of  $P$ :

**Lemma 2.3.** *Let  $E$  be an edge of  $P$  with primitive inner normal vector  $w \in M$ . Then  $P$  admits a mutation with respect to  $w$  if and only if  $|E \cap N| - 1 \geq r_E$ .*

*Proof.* Let  $k := |E \cap N| - 1$  be the length of  $E$ . At height  $h = h_{\min} = -r_E$ , condition (2.1) becomes  $E = G_{h_{\min}} + r_E F$ . Hence this condition can be satisfied if and only if  $k \geq r_E$ . Suppose that  $k \geq r_E$  and consider the cone  $C := \text{cone}(E)$  generated by  $E$ . At height  $h_{\min} < h < 0$ ,  $h \in \mathbb{Z}$ , the line segment  $C_h := \{v \in C \mid w(v) = h\} \subset N_{\mathbb{Q}}$  (with rational end-points) has length  $|h|k/r_E \geq |h|$ . Hence  $w_h(C) \subset w_h(P)$  has length at least  $|h| - 1$ . Suppose that there exists some  $v \in \mathcal{V}(P)$  such that  $w(v) = h$ . Since  $v \notin w_h(C)$  we conclude that  $w_h(P)$  has length at least  $|h|$ . Hence condition (2.1) can be satisfied. If  $\{v \in \mathcal{V}(P) \mid w(v) = h\} = \emptyset$  then we can simply take  $G_h = \emptyset$  to satisfy condition (2.1).  $\square$

Lemma 2.3 states that the existence of a mutation of a polygon is completely determined by a purely local condition on edge length (this is not the case in higher dimensions). As a consequence, we can rephrase mutation of polygons as follows.

**Corollary 2.4** (c.f. [1]). *Choose an orientation of  $N$  and label the vertices of  $P$  by  $v_1, v_2, \dots$  counterclockwise, such that  $w(v_1) = h_{\max}$  and  $w(v_2) \neq h_{\max}$ . Let  $f \in w^{\perp} \cap N$  be a primitive lattice element. Then  $P$  admits a mutation with respect to  $w$  and the unit line segment  $F = \text{conv}\{\mathbf{0}, f\}$  if and only if there exists an edge  $\text{conv}\{v_i, v_{i+1}\}$  such that  $w(v_i) = w(v_{i+1}) = h_{\min}$  and  $v_{i+1} - v_i = kf$ , where  $k \geq -h_{\min}$  is a positive integer. In order to describe the mutation, we distinguish between two cases:*

- (I)  $P$  has  $n$  vertices  $v_1, \dots, v_n$  and  $v_1$  is the unique maximum for  $w$  on  $P$ ;
- (II)  $P$  has  $n + 1$  vertices  $v_1, \dots, v_{n+1}$  and  $w(v_1) = w(v_{n+1}) = h_{\max}$ .

Then the mutation of  $P$  with respect to  $w$  is the Fano polygon with  $n + 1$  vertices  $v'_1, \dots, v'_{n+1}$  given by:

Case (I)	Case (II)
$v'_j = \begin{cases} v_j, & \text{if } 1 \leq j \leq i; \\ v_j + w(v_j)f, & \text{if } i < j \leq n; \\ v_1 + h_{\max}f, & \text{if } j = n + 1. \end{cases}$	$v'_j = \begin{cases} v_j, & \text{if } 1 \leq j \leq i; \\ v_j + w(v_j)f, & \text{if } i < j \leq n; \\ v_{n+1} + h_{\max}f, & \text{if } j = n + 1. \end{cases}$

2.2. **Mutation in  $M$ .** Given a Fano polygon  $P \subset N_{\mathbb{Q}}$  we define the *dual polygon*

$$P^* := \{u \in M_{\mathbb{Q}} \mid u(v) \geq -1 \text{ for all } v \in P\} \subset M_{\mathbb{Q}}.$$

In general this has rational-valued vertices and necessarily contains the origin in its strict interior. Define  $\varphi : M_{\mathbb{Q}} \rightarrow M_{\mathbb{Q}}$  by  $u \mapsto u - u_{\min} w$ , where  $u_{\min} := \min\{u(v) \mid v \in F\}$ . Since  $F = \text{conv}\{\mathbf{0}, v_E\}$ , this is equivalent to


$$\varphi(u) = \begin{cases} u, & \text{if } u(v_E) \geq 0; \\ u - u(v_E)w, & \text{if } u(v_E) < 0. \end{cases}$$

This is a piecewise- $GL_2(\mathbb{Z})$  map, partitioning  $M_{\mathbb{Q}}$  into two half-spaces whose common boundary is generated by  $w$ . Crucially [2, Proposition 4 and the discussion on pg. 12]:

$$\varphi(P^*) = Q^*, \quad \text{where } Q := \text{mut}_w(P, F).$$

An immediate consequence of this is that the area and Ehrhart series of the dual polygons are preserved under mutation:  $\text{Vol}(P^*) = \text{Vol}(Q^*)$  and  $\text{Ehr}_{P^*}(t) = \text{Ehr}_{Q^*}(t)$ . Equivalently, mutation preserves the anti-canonical degree and Hilbert series of the corresponding toric varieties:  $(-K_{X_P})^2 = (-K_{X_Q})^2$  and  $\text{Hilb}(X_P, -K_{X_P}) = \text{Hilb}(X_Q, -K_{X_Q})$ .


**Example 2.5.** Consider the polygon  $P_{(1,1,1)} := \text{conv}\{(1,1), (0,1), (-1,-2)\} \subset N_{\mathbb{Q}}$ . The toric variety corresponding to  $P_{(1,1,1)}$  is  $\mathbb{P}^2$ . Let  $w = (0,-1) \in M$ , so that  $h_{\min} = -1$  and  $h_{\max} = 2$ , and set  $F = \text{conv}\{\mathbf{0}, (1,0)\} \subset N_{\mathbb{Q}}$ . Then  $F$  is a factor of  $P_{(1,1,1)}$  with respect to  $w$ , giving the mutation  $P_{(1,1,2)} := \text{mut}_w(P_{(1,1,1)}, F)$  with vertices  $(0,1), (-1,-2), (1,-2)$  as depicted below. The toric variety corresponding to  $P_{(1,1,2)}$  is  $\mathbb{P}(1,1,4)$ .




In  $M_{\mathbb{Q}}$  we see the mutation as a piecewise- $GL_2(\mathbb{Z})$  transformation. This acts on the left-hand half-space  $\{(u_1, u_2) \in M_{\mathbb{Q}} \mid u_1 < 0\}$  via the transformation

$$(u_1, u_2) \mapsto (u_1, u_2) \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$$

and on the right-hand half-space via the identity.



We can draw a graph of all possible mutations obtainable from  $P_{(1,1,1)}$ : the vertices of the graph denote  $GL_2(\mathbb{Z})$ -equivalence classes of Fano polygons, and two vertices are connected by an edge if there exists a mutation between the two Fano polygons (notice that, since mutations are invertible, we can regard the edges as being undirected). We obtain a tree whose typical vertex is trivalent [4, Example 3.14]:



Here the vertices have been labelled with weights  $(a, b, c)$ , where the polygon  $P_{(a,b,c)}$  corresponds to the toric variety  $\mathbb{P}(a^2, b^2, c^2)$ . The triples  $(a, b, c)$  are solutions to the Markov equation

$$3xyz = x^2 + y^2 + z^2,$$

and each mutation corresponds, up to permutation of  $a, b$ , and  $c$ , to a transformation of the form  $(a, b, c) \mapsto (3bc - a, b, c)$ . In the theory of Markov equations these transformations are also called *mutations*. A solution  $(a, b, c)$  is called *minimal* if  $a + b + c$  is minimal, and every solution can be reached via mutation from a minimal solution. Minimal solutions correspond to those triangles with  $\text{Vol}(P_{(a,b,c)})$  minimal. In this example  $(1, 1, 1)$  is the unique minimal solution. These statements can be generalised to any mutation between triangles [4]. Hacking–Prokhorov [25] use these minimal solutions in their classification of rank-one qG-smoothable del Pezzo surfaces of class TG.

### 3. INVARIANTS OF FANO POLYGONS

We wish to be able to establish whether or not two Fano polygons are mutation-equivalent. In this section we introduce two mutation invariants of a Fano polygon  $P \subset N_{\mathbb{Q}}$ : singularity content, discussed in §3.1 below, can be thought of as studying the part of  $P$  that remains untouched by mutation (the basket  $\mathcal{B}$  of residual singularities); the cluster algebra  $\mathcal{A}_P$ , discussed in §3.3 below, studies the part of  $P$  that changes under mutation (the primitive  $T$ -singularities). Although we have no proof, it seems likely that together these two invariants completely characterise the mutation-equivalence classes. Finally, in §3.4 we briefly mention the connection with affine manifolds and the Gross–Seibert program [22].

**3.1. Singularity content.** In [3] the concept of *singularity content* for a Fano polygon was introduced. First we state the definition for a cyclic quotient singularity  $\frac{1}{R}(a, b)$ , where  $\gcd\{R, a\} = \gcd\{R, b\} = 1$ . (Recall that  $\frac{1}{R}(a, b)$  denotes the germ of a quotient singularity  $\mathbb{C}^2/\mu_R$ , where  $\varepsilon \in \mu_R$  acts via  $(x, y) \mapsto (\varepsilon^a x, \varepsilon^b y)$ .) Let  $k, r, c \in \mathbb{Z}$  be non-negative integers such that  $k = \gcd\{R, a + b\}$ ,  $R = kr$ , and  $a + b = kc$ . Then  $r$  is equal to the *Gorenstein index* of the singularity, and  $k$  is called the *length* (or *width*). Thus  $\frac{1}{R}(a, b)$  can be written in the form  $\frac{1}{kr}(1, kc - 1)$  for some  $c \in \mathbb{Z}$  with  $\gcd\{r, c\} = 1$ .

**Definition 3.1** ( $r \mid k$ ). A cyclic quotient singularity such that  $k = nr$  for some  $n \in \mathbb{Z}_{>0}$ , that is, a cyclic quotient singularity of the form  $\frac{1}{nr^2}(1, nrc - 1)$ , is called a *T-singularity* or a singularity of *class T*. When  $n = 1$ , so that the singularity is of the form  $\frac{1}{r^2}(1, rc - 1)$ , we call it a *primitive T-singularity*.

**Definition 3.2** ( $k < r$ ). A cyclic quotient singularity of the form  $\frac{1}{kr}(1, kc - 1)$  with  $k < r$  is called a *residual singularity* or a singularity of *class R*.

$T$ -singularities appear in the work of Wahl [41] and Kollár–Shepherd-Barron [29]. A cyclic quotient singularity is of class  $T$  if and only if it admits a qG-smoothing. At the opposite extreme, a singularity is of class  $R$  if and only if it is rigid under qG-deformation. More generally, consider the cyclic quotient singularity  $\sigma = \frac{1}{kr}(1, kc - 1)$ . Let  $0 \leq k_0 < r$  and  $n$  be the unique non-negative integers such that  $k = nr + k_0$ . Then either  $k_0 = 0$  and  $\sigma$  is qG-smoothable, or  $k_0 > 0$  and  $\sigma$  admits a qG-deformation to the residual singularity  $\frac{1}{k_0r}(1, k_0c - 1)$  [1, 3]. This motivates the following definition:

**Definition 3.3** ([3, Definition 2.4]). With notation as above, let  $\sigma = \frac{1}{kr}(1, kc - 1)$  be a cyclic quotient singularity. The *residue* of  $\sigma$  is given by

$$\text{res}(\sigma) := \begin{cases} \emptyset & \text{if } k_0 = 0, \\ \frac{1}{k_0r}(1, k_0c - 1) & \text{otherwise.} \end{cases}$$

The *singularity content* of  $\sigma$  is given by the pair  $\text{SC}(\sigma) := (n, \text{res}(\sigma))$ .

**Example 3.4.** Let  $\sigma = \frac{1}{nr^2}(1, nrc - 1)$  be a  $T$ -singularity. Then  $\text{SC}(\sigma) = (n, \emptyset)$ .

Singularity content has a natural description in terms of the cone  $C$  defining the singularity. We call a two-dimensional cone  $C \subset N_{\mathbb{Q}}$  a *T-cone* (respectively *primitive T-cone*) if the corresponding cyclic quotient singularity is a  $T$ -singularity (respectively primitive  $T$ -singularity), and we call  $C$  an *R-cone* if the corresponding singularity is a residual singularity. Let  $C = \text{cone}\{\rho_0, \rho_1\} \subset N_{\mathbb{Q}}$  be a

two-dimensional cone with rays generated by the primitive lattice points  $\rho_0$  and  $\rho_1$  in  $N$ . The line segment  $E = \text{conv}\{\rho_0, \rho_1\}$  is at height  $r$  and has length  $|E \cap N| - 1 = k$ . Write  $k = nr + k_0$ . Then there exists a partial crepant subdivision of  $C$  into  $n$  cones  $C_1, \dots, C_n$  of length  $r$  and, if  $k_0 \neq 0$ , one cone  $C_0$  of length  $k_0 < r$ . Although not immediately obvious from this description, the singularity corresponding to the  $R$ -cone  $C_0$  is well-defined and equal to  $\frac{1}{k_0 r}(1, k_0 c - 1)$ , where we refer to [3] for the precise definition of  $c$ . The singularities corresponding to the  $n$  primitive  $T$ -cones  $C_1, \dots, C_n$  depend upon the particular choice of subdivision: again, see [3, Proposition 2.3] for the precise statement.

**Definition 3.5.** Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon with edges  $E_1, \dots, E_m$ , numbered cyclically, and let  $\sigma_1, \dots, \sigma_m$  be the corresponding two-dimensional cyclic quotient singularities  $\sigma_i = \text{cone}(E_i)$ , with  $\text{SC}(\sigma_i) = (n_i, \text{res}(\sigma_i))$ . The *singularity content* of  $P$ , denoted by  $\text{SC}(P)$ , is the pair  $(n, \mathcal{B})$ , where  $n := \sum_{i=1}^m n_i$  and  $\mathcal{B}$  is the cyclically-ordered list  $\{\text{res}(\sigma_i) \mid 1 \leq i \leq m, \text{res}(\sigma_i) \neq \emptyset\}$ . We call  $\mathcal{B}$  the *basket* of residual singularities of  $P$ .

Singularity content is a mutation invariant of  $P$  [3, Proposition 3.6]. Intuitively one can see this from Lemma 2.3: mutation removes a line segment of length  $|h_{\min}|$  from the edge at height  $h_{\min}$ , changing the corresponding singularity content by  $(n, \text{res}(\sigma)) \mapsto (n - 1, \text{res}(\sigma))$ ; mutation adds a line segment of length  $h_{\max}$  at height  $h_{\max}$ , changing the singularity content by  $(n', \text{res}(\sigma')) \mapsto (n' + 1, \text{res}(\sigma'))$ . Put another way, mutation removes a primitive  $T$ -cone with Gorenstein index  $-h_{\min}$  and adds a primitive  $T$ -cone with Gorenstein index  $h_{\max}$ , leaving the residual cones unchanged. We can rephrase Lemma 2.3 in terms of singularity content:

**Lemma 3.6.** *Let  $E$  be an edge of  $P$  with primitive inner normal vector  $w \in M$ , and let  $(n, \text{res}(\sigma))$  be the singularity content of  $\sigma = \text{cone}(E)$ . Then  $P$  admits a mutation with respect to  $w$  if and only if  $n \neq 0$ .*

Singularity content provides an upper-bound on the maximum number of vertices of any polygon  $P$  with  $\text{SC}(P) = (n, \mathcal{B})$ , or, equivalently, an upper-bound on the Picard rank  $\rho$  of the corresponding toric variety  $X_P$  [3, Lemma 3.8]:

$$(3.1) \quad |\mathcal{V}(P)| \leq n + |\mathcal{B}|, \quad \rho \leq n + |\mathcal{B}| - 2.$$

**Example 3.7.** Consider  $P_{(1,1)} = \text{conv}\{(1,0), (0,1), (-1,-3)\}$  with corresponding toric variety  $\mathbb{P}(1,1,3)$ . This has singularity content  $(2, \{\frac{1}{3}(1,1)\})$ , hence, by (3.1), any polygon  $Q$  mutation-equivalent to  $P_{(1,1)}$  has three vertices. Mutations between triangles were characterised in [4]: the mutation graph is given by

$$(1,1) \text{ ——— } (1,4) \text{ ——— } (4,19) \text{ ——— } (19,91) \text{ ——— } (91,436) \text{ ——— } \dots$$

Here the vertices have been labelled by pairs  $(a,b) \in \mathbb{Z}_{>0}^2$ , and correspond to  $\mathbb{P}(a^2, b^2, 3)$  and its associated triangle. These pairs are solutions to the Diophantine equation  $5xy = x^2 + y^2 + 3$ . Up to exchanging  $a$  and  $b$ , a mutation of triangles corresponds to the mutation  $(a,b) \mapsto (5b - a, b)$  of solutions. There is a unique minimal solution given by  $(1,1)$ .

As noted in §1.1, the toric variety  $X_P$  is qG-deformation-equivalent to a del Pezzo surface  $X$  with singular points  $\mathcal{B}$  and topological Euler number of  $X \setminus \text{Sing}(X)$  equal to  $n$  [1, 3]. The degree and Hilbert series can be expressed purely in terms of singularity content. Recall that information about a minimal resolution of a singularity  $\sigma = \frac{1}{R}(1, a - 1)$  is encoded in the Hirzebruch–Jung continued fraction expansion  $[b_1, \dots, b_s]$  of  $R/(a - 1)$ ; see, for example, [18]. For each  $i \in \{1, \dots, s\}$  we inductively define the positive integers  $\alpha_i, \beta_i$  as follows:

$$\begin{aligned} \alpha_1 &= \beta_s = 1, \\ \alpha_i / \alpha_{i-1} &:= [b_{i-1}, \dots, b_1], \quad 2 \leq i \leq s, \\ \beta_i / \beta_{i+1} &:= [b_{i+1}, \dots, b_s], \quad 1 \leq i \leq s - 1. \end{aligned}$$

The values  $-b_i$  give the self-intersection numbers of the exceptional divisors of the minimal resolution of  $\sigma$ , and the values  $d_i := -1 + (\alpha_i + \beta_i)/R$  give the discrepancies. The degree contribution of  $\sigma$  is given by:

$$A_{\sigma} := s + 1 - \sum_{i=1}^s d_i^2 b_i + 2 \sum_{i=1}^{s-1} d_i d_{i+1}.$$



The Riemann–Roch contribution  $Q_\sigma$  of  $\sigma$  can be computed in terms of Dedekind sums (see [38, §8]):

$$Q_\sigma = \frac{1}{1-t^R} \sum_{i=1}^{R-1} (\delta_{ai} - \delta_0) t^{i-1}, \quad \text{where} \quad \delta_j := \frac{1}{R} \sum_{\substack{\varepsilon \in \mu_R \\ \varepsilon \neq 1}} \frac{\varepsilon^j}{(1-\varepsilon)(1-\varepsilon^{a-1})}.$$

**Proposition 3.8** ([3, Proposition 3.3 and Corollary 3.5]). *Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon with singularity content  $(n, \mathcal{B})$ . Let  $X_P$  be the toric variety given by the spanning fan of  $P$ . Then*

$$(-K_{X_P})^2 = 12 - n - \sum_{\sigma \in \mathcal{B}} A_\sigma \quad \text{and} \quad \text{Hilb}(X_P, -K_{X_P}) = \frac{1 + ((-K_{X_P})^2 - 2)t + t^2}{(1-t)^3} + \sum_{\sigma \in \mathcal{B}} Q_\sigma.$$

The terms  $Q_\sigma$  can be interpreted as a periodic correction to the initial term

$$\frac{1 + ((-K_{X_P})^2 - 2)t + t^2}{(1-t)^3} = \sum_{i \geq 0} \binom{i+1}{2} (-K_{X_P})^2 + 1 \Big) t^i.$$

Set  $Q_{\text{num}} := (1-t^R)Q_\sigma$ . The contribution from  $Q_\sigma$  at degree  $i$  is equal to the coefficient of  $t^m$  in  $Q_{\text{num}}$ , where  $i \equiv m \pmod{R}$ .

**Example 3.9.** Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon with singularity content  $(n, \{m \times \frac{1}{3}(1, 1)\})$ , for some  $n \in \mathbb{Z}_{\geq 0}$ ,  $m \in \mathbb{Z}_{>0}$ . We see that  $A_{\frac{1}{3}(1,1)} = 5/3$  and  $Q_{\frac{1}{3}(1,1)} = -t/3(1-t^3)$ , giving

$$\begin{aligned} \text{Vol}(P^*) &= (-K_{X_P})^2 = 12 - n - \frac{5m}{3} \quad \text{and} \\ \text{Ehr}_{P^*}(t) &= \text{Hilb}(X_P, -K_{X_P}) = \frac{1 + (11 - n - 2m)t + (12 - n - m)t^2 + (11 - n - 2m)t^3 + t^4}{(1-t^3)(1-t)^2}. \end{aligned}$$

In particular, for any  $i \geq 0$ ,

$$|iP^* \cap M| = h^0(X_P, -iK_{X_P}) = \binom{i+1}{2} (-K_{X_P})^2 + 1 - \begin{cases} \frac{m}{3} & \text{if } i \equiv 1 \pmod{3}, \\ 0 & \text{otherwise.} \end{cases}$$

Since  $|P^* \cap M| = 13 - n - 2m \geq 1$  we have that  $0 \leq n \leq 10$  and  $1 \leq m \leq 6 - n/2$ . Notice that  $(5, \{4 \times \frac{1}{3}(1, 1)\})$ ,  $(3, \{5 \times \frac{1}{3}(1, 1)\})$ , and  $(1, \{6 \times \frac{1}{3}(1, 1)\})$  give  $h^0(X, -K_X) = 0$ , hence there cannot exist a corresponding Fano polygon  $P$  (or toric surface  $X_P$ ). They do, however, correspond to the Euler numbers and singular points of the del Pezzo surfaces  $X_{4,1/3}$ ,  $X_{5,2/3}$ , and  $X_{6,7/2}$ , respectively, described in [14]. These three del Pezzo surfaces cannot be of class TG. It is tempting to conjecture that having  $h^0(X, -K_X) = 0$  is the only obstruction to a del Pezzo surface  $X$  being of class TG.

**3.2. The sublattice  $\Gamma_P$  of  $M$ .** There exist examples of Fano polygons that have the same singularity content, but that are not mutation-equivalent (see Example 3.12 below); there is additional structure in the arrangement of the primitive  $T$ -singularities that singularity content ignores.

**Definition 3.10.** Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon. Define  $\Gamma_P$  to be the sublattice of  $M$  generated by all primitive inner normal vectors  $w$  to an edge  $E$  of  $P$  with  $|E \cap N| - 1 \geq r_E$ . Let  $[M : \Gamma_P]$  denote the index of this sublattice in  $M$ .<sup>1</sup>

**Lemma 3.11.** *Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon and let  $Q := \text{mut}_w(P, F)$  be a mutation of  $P$ . Then  $[M : \Gamma_P] = [M : \Gamma_Q]$ .*

*Proof.* Recall from §2.2 that mutation acts on the element of  $M$  via the piecewise-linear map  $\varphi : u \mapsto u - u_{\min} w$ , where  $u_{\min} := \min\{u(v_F) \mid v_F \in \mathcal{V}(F)\}$ . By Lemma 2.3,  $w \in \Gamma_P$  and  $\varphi$  maps  $\Gamma_P$  to itself. Mutations are invertible, with the inverse to  $\text{mut}_w(\cdot, F)$  being  $\text{mut}_{-w}(\cdot, F)$ . Thus the index is preserved.  $\square$

<sup>1</sup>We thank one of the referees for making the following observation. After making a generic partial smoothing  $X$  of  $X_P$  the group  $M/\Gamma_P$  is the fundamental group of the complement  $U$  of an anti-canonical divisor in  $X$ . Indeed,  $U$  is homotopy equivalent to a torus with discs glued along representatives of homology classes defined by vanishing cycles of a (singular) torus fibration from  $U$  to a disc, and these classes are precisely the vectors  $w \in M$  in Definition 3.10.

**Example 3.12.** Consider the reflexive polygons

$$R_{15} := \text{conv}\{(1, 0), (0, 1), (-1, -1), (0, -1)\} \subset N_{\mathbb{Q}}$$

$$\text{and } R_{16} := \text{conv}\{(1, 0), (0, 1), (-1, 0), (0, -1)\} \subset N_{\mathbb{Q}}.$$

The toric varieties defined via the spanning fan are the del Pezzo surfaces  $\mathbb{F}_1$  and  $\mathbb{P}^1 \times \mathbb{P}^1$  respectively. Both  $R_{15}$  and  $R_{16}$  have singularity content  $(4, \emptyset)$ . The primitive normal vectors to the edges of  $R_{15}$  generate all of  $M$ , whereas the normal vectors of the edges of  $R_{16}$  generate an index two sublattice  $\Gamma_P = \langle (1, 1), (-1, 1) \rangle \subset M$ . Lemma 3.11 shows that  $R_{15}$  and  $R_{16}$  are not mutation-equivalent. We will give another proof of this, using quiver mutations, in Example 3.19 below.

**3.3. Quivers and cluster algebras.** We first recall the definition of cluster algebra [16, 17], having fixed a rank- $n$  lattice  $L$  and a skew-symmetric form  $\{\cdot, \cdot\}$ .

**Definition 3.13.** A *seed* is a pair  $(B, C)$  where  $B$  is a basis of  $L$  and  $C$  is a transcendence basis of  $\mathbb{C}(L)$ , referred to as a *cluster*.

**Definition 3.14.** Given a seed  $(B, C)$  with  $B = \{e_1, \dots, e_n\}$  and  $C = \{x_1, \dots, x_n\}$ , the  $j$ th mutation of  $(B, C)$  is the seed  $(B', C')$ , where  $B' = \{e'_1, \dots, e'_n\}$  and  $C' = \{x'_1, \dots, x'_n\}$  are defined by:

$$e'_k = \begin{cases} -e_j, & \text{if } k = j \\ e_k + \max(b_{kj}, 0)e_j, & \text{otherwise} \end{cases}$$

where  $b_{kj} = \{e_k, e_j\}$ ,

$$x'_k = x_k \text{ if } k \neq j, \quad \text{and} \quad x_j x'_j = \prod_{\substack{k \text{ such that} \\ b_{jk} > 0}} x_k^{b_{jk}} + \prod_{\substack{l \text{ such that} \\ b_{jl} < 0}} x_l^{b_{lj}}.$$

Recall from §1.2 that there is a quiver  $Q_P$  and a cluster algebra  $\mathcal{A}_P$  associated to a Fano polygon  $P$ . Let  $\text{SC}(P) = (n, \mathcal{B})$  and fix a numbering of the  $n$  primitive  $T$ -cones in the spanning fan of  $P$ . The  $i$ th primitive  $T$ -cone in the spanning fan of  $P$  corresponds to a vertex  $v_i$  of  $Q_P$ , and thus corresponds to a basis element  $e_i$  in  $B_P$ , where  $(B_P, C_P)$  is the initial seed for  $\mathcal{A}_P$ . Let  $E_i$  denote the edge of  $P$  determined by the  $i$ th primitive  $T$ -cone; note that different primitive  $T$ -cones can determine the same edge.

**Proposition 3.15** (Mutations of seeds induce mutations of polygons). *Let  $P$  be a Fano polygon with singularity content  $(n, \mathcal{B})$ . Denote by  $w_i \in M$  the primitive inner normal vector to  $E_i$ . Consider the map  $\pi: L \rightarrow M$  such that  $\pi(e_i) = w_i$  for each  $i$ . Let  $(B_P, C_P)$  be the initial seed for  $\mathcal{A}_P$ , and write  $B = \{e_1, \dots, e_n\}$ ,  $C = \{x_1, \dots, x_n\}$ . Let  $(B', C')$  be the  $j$ th mutation of  $(B_P, C_P)$ , and write  $B' = \{e'_1, \dots, e'_n\}$ . Then  $\{\pi(e'_1), \dots, \pi(e'_n)\}$  are the primitive inner normal vectors to the edges of  $P'$ , where  $P'$  is the mutation of  $P$  determined by  $w_j$ . Furthermore, every mutation  $P'$  of  $P$  arises in this way.*

*Proof.* This is a straightforward calculation using mutation in  $M$  (see §2.2).  $\square$

There is a well-known notion of quiver mutation, going back to Bernstein–Gelfand–Ponomarev [9], Fomin–Zelevinsky [17], and others.

**Definition 3.16.** Given a quiver  $Q$  and a vertex  $v$  of  $Q$ , the *mutation of  $Q$  at  $v$*  is the quiver  $\text{mut}(Q, v)$  obtained from  $Q$  by:

- (i) adding, for each subquiver  $v_1 \rightarrow v \rightarrow v_2$ , an arrow from  $v_1$  to  $v_2$ ;
- (ii) deleting a maximal set of disjoint two-cycles;
- (iii) reversing all arrows incident to  $v$ .

The resulting quiver is well-defined up to isomorphism, regardless of the choice of two-cycles in (ii).

**Proposition 3.17** (Mutations of polygons induce mutations of quivers). *Let  $P$  be a Fano polygon, let  $v$  be a vertex of  $Q_P$  corresponding to a primitive  $T$ -cone in  $P$ , and let  $P'$  be the corresponding mutation of  $P$ . We have  $Q_{P'} = \text{mut}(Q_P, v)$ .*

*Proof.* Let  $E$  denote the edge of  $P$  determined by the primitive  $T$ -cone corresponding to  $v$ , and let  $w \in M$  denote the primitive inner normal vector to  $E$ . Mutation with respect to  $w$  acts on  $M$  as a piecewise-linear transformation that is the identity in one half-space, and on the other half-space is a shear transformation  $u \mapsto u + (w \wedge u)w$ . Thus determinants between the pairs of normal vectors change as follows:

- (i) The inner normal vector  $w$  to the mutating edge  $E$  becomes  $-w$ , so that all arrows into  $v$  change direction;
- (ii) For a pair of normal vectors in the same half-space (as defined by  $w$ ), the determinant does not change;
- (iii) Consider primitive  $T$ -cones with inner normal vectors in different half-spaces (as defined by  $w$ ), let the corresponding vertices of  $Q_P$  be  $v_1$  and  $v_2$ , and let the corresponding inner normal vectors in  $M$  be  $w_1$  and  $w_2$ . Without loss of generality we may assume that  $w_1 \wedge w > 0$  and  $w_2 \wedge w < 0$ , so that there are arrows  $v_1 \rightarrow v \rightarrow v_2$  in  $Q_P$ . Under mutation, the primitive inner normal vectors change as  $w_1 \mapsto w'_1$ ,  $w_2 \mapsto w'_2$  where  $w'_1 = w_1$ ,  $w'_2 = w_2 + (w \wedge w_2)w$ . Thus:

$$w'_1 \wedge w'_2 = w_1 \wedge w_2 + (w \wedge w_2)(w_1 \wedge w)$$

and so we add an arrow for each path  $v_1 \rightarrow v \rightarrow v_2$ . Cancelling two-cycles results in precisely the result of calculating the signed total number of arrows from  $v_1$  to  $v_2$ .


Observing finally that if  $v_1, v_2$  give normal vectors in the same half-space then there are no paths  $v_1 \rightarrow v \rightarrow v_2$  or  $v_2 \rightarrow v \rightarrow v_1$ , we see that this description coincides with that of a quiver mutation.  $\square$

Propositions 3.15 and 3.17 give upper and lower bounds on the mutation graphs of Fano polygons. For example:

**Corollary 3.18** (see Example 1.4). *If a Fano polygon  $P$  has singularity content  $(2, \mathcal{B})$  and is such that the primitive inner normal vectors of the two edges corresponding to the two primitive  $T$ -cones form a basis of the dual lattice  $M$  (that is, if  $\Gamma_P = M$ ), then the mutation-equivalence class of  $P$  has at most five members.*

*Proof.* The quiver associated to  $P$  is simply the  $A_2$  quiver. The cluster algebra  $\mathcal{A}_P$  is therefore well-known (being the cluster algebra associated to the  $A_2$  quiver) and its cluster exchange graph forms a pentagon. Note however that the *quiver* mutation graph is trivial, as the  $A_2$  quiver mutates only to itself. Proposition 3.15 implies that the mutation graph of  $P$  has at most five vertices. (Proposition 3.17 does not give a non-trivial lower bound here: indeed polygon 7 in Figure 2 on page 27 gives an example of such a polygon  $P$  with trivial mutation graph.)  $\square$

**Example 3.19.** For the polygons  $R_{15}$  and  $R_{16}$  considered in Example 3.12 above the associated quivers  $Q_{R_{15}}$  and  $Q_{R_{16}}$  are:



Observe that for  $Q_{R_{16}}$  the number of arrows between any two vertices is even. It is easy to see that this property is preserved under mutation. Therefore the quivers  $Q_{R_{15}}$  and  $Q_{R_{16}}$  are not mutation-equivalent, and so the polygons  $R_{15}$  and  $R_{16}$  are not mutation-equivalent.

**Remark 3.20.** Following Fock–Goncharov [16], Gross–Hacking–Keel [20], and Mandel [33], there is a rich interplay between the theory of cluster algebras and the geometry of log Calabi–Yau surfaces.

In [20, Construction 5.3] a cluster structure analogous to the one defined in §1.2 is given. In this construction, the collection of basis vectors  $e_i$  comprising a seed is mapped to a collection of vectors  $w_i$  which are interpreted as generating elements for the rays of a fan in a two dimensional lattice. This toric surface is then blown up at  $k$  points on each divisor  $D_w$ , where  $k$  is the number of seed vectors mapping to  $w$ , and the family of such surfaces is denoted  $\mathcal{Y}$ . Combining this with the construction described in §1.2, given a Fano polygon  $P$  we can produce a log Calabi–Yau surface by blowing up

$m$  points on the boundary of a toric surface given by the normal fan to  $P$ , where  $m$  is the singularity content of the corresponding edge of  $P$ .

Following [20] further, the family of surfaces  $\mathcal{Y}$  obtained by this elementary birational construction agrees, up to codimension two, with the ‘ $\mathcal{X}$ -type cluster variety’ defined in [16]. Indeed, Fock–Goncharov define a pair of (mirror-dual) varieties (the  $\mathcal{X}$ - and  $\mathcal{A}$ -type cluster varieties) directly from the data of a cluster algebra. Interpreting  $\mathcal{Y}$  as a cluster variety, it is covered by *seed tori* whose transition functions are the mutations appearing in [2].

The expected mirror-duality in this context is between the  $\mathcal{X}$ -type cluster variety equipped with a superpotential  $W$ , and the generic smoothing of the toric variety  $X_P$ . Indeed, a mirror construction for log Calabi–Yau surfaces is described by Gross–Hacking–Keel in [21], and in [36] it is shown that, in this context, the corresponding family does indeed arise from the smoothing of the toric variety  $X_P$ .

**3.4. Affine manifolds.** This section sketches a more geometric approach to finding mutation invariants of polygons, based on the foundational papers of Gross–Siebert [23] and Kontsevich–Soibelman [30]. In broad terms, one wishes to generalise the fact that the dual polygon  $P^*$  is the base of a special Lagrangian torus fibration given by the moment map, by allowing more general bases and more general torus fibrations. Specifically, the base of a special Lagrangian torus fibration carries an *affine structure* [22, 30]. Deforming the torus fibration then leads to a deformation of the “candidate base” (actually exhibiting a fibration over the deformed base is fraught with difficulty); the deformed base does not have the structure of a polygon, but rather is an *affine manifold*.

**Remark 3.21.** Whilst the passage from affine manifolds to algebraic geometry requires sophisticated technology from the programs of Kontsevich–Soibelman and Gross–Siebert, a construction of a *Lagrangian* fibration over this affine manifold with “smoothed corners” was described by Symington [40].

**Definition 3.22.** A two-dimensional *integral affine manifold*  $B$  is a manifold which admits a maximal atlas with transition functions in  $GL_2(\mathbb{Z}) \times \mathbb{Z}^2$

**Example 3.23.** The interior of  $P^*$  is an example of an integral affine manifold, covered as it is by a single chart. The polygon  $P^*$  itself is an example of an *affine manifold with corners*.

Allowing singular fibres in the special Lagrangian torus fibration corresponds roughly to the base manifold acquiring *focus-focus singularities*. We say “roughly” because, following Gross–Siebert [22], we should be considering toric *degenerations* rather than torus fibrations; see [23, 30]. The local model for such a singularity is  $\mathbb{R}^2 \setminus \{0\}$ , regarded as an affine manifold via a cover with two charts

$$U_1 = \mathbb{R}^2 \setminus \mathbb{R}_{\leq 0} \times \{0\} \quad \text{and} \quad U_2 = \mathbb{R}^2 \setminus \mathbb{R}_{\geq 0} \times \{0\}$$

with the transition function

$$(x, y) \mapsto \begin{cases} (x, y), & \text{if } y > 0, \\ (x + y, y), & \text{if } y < 0. \end{cases}$$

Sliding singularities, which Kontsevich–Soibelman call *moving worms* [30], gives an affine analogue of the deformations of the varieties. In particular, allowing singularities to collide with boundary points of the affine manifold creates corners and provides an analogue for the toric qG-degenerations of the surface. This process, and its lifting via the Gross–Siebert program to construct the corresponding degeneration of algebraic varieties, are described in [36].

Given the dual polygon  $P^*$ , the set of singularities that we can introduce is in natural bijection with the primitive  $T$ -cones appearing in the singularity content of  $P$ . Consider the following process. Take a primitive  $T$ -cone  $\sigma$  of  $P$ , and introduce the corresponding singularity into the interior of  $P^*$ , partially smoothing that corner and forming an affine manifold  $B$ . Now slide this singularity along the monodromy-invariant line, all the way to the opposite of  $B$ , and so forming a polygon  $P'^*$  with dual polygon  $P'$ .

**Lemma 3.24.**  $P'$  is equal to the mutation of  $P$  defined by the chosen primitive  $T$ -cone  $\sigma$ .

*Proof.* Mutation induces a piecewise-linear transformation on the dual polygon. In fact this corresponds exactly to the transition function between the two charts defining  $B$ . As the singularity

approaches a corner, one of the charts covers all but a line segment with vanishing length, hence the polygons are related by the piecewise-linear transition function applied to the entire polygon.  $\square$

Given a polygon  $P^*$  one can introduce a maximal set of singularities and thus form an affine manifold  $B$ . Regarding two affine manifolds which differ by moving singularities along monodromy-invariant lines as equivalent, we see that every polygon in the same mutation class is equivalent as an affine manifold. This gives a mutation invariant – the affine manifold  $B$  – which we can use to distinguish minimal polygons. Specifically, we can compare the respective *monodromy representations*. Since the transition functions of  $B$  are in  $GL_2(\mathbb{Z})$  we can define parallel transport of integral vector fields. Fixing a basepoint in  $B$ , parallel transport around loops gives a representation  $\pi_1(B) \rightarrow SL_2(\mathbb{Z})$ . Properties of this representation may be used to distinguish different mutation classes of polygons. For example, in the cases considered in Example 3.12, one monodromy representation is surjective and the other is not.

#### 4. MINIMAL FANO POLYGONS

Given a polygon  $P \subset N_{\mathbb{Q}}$  we want to find a preferred representative in the mutation-equivalence class of  $P$ . Let  $\partial P$  denote the boundary of  $P$  and let  $P^\circ := P \setminus \partial P$  denote the strict interior of  $P$ . We introduce the following definition:

**Definition 4.1.** We call a Fano polygon  $P \subset N_{\mathbb{Q}}$  *minimal* if for every mutation  $Q := \text{mut}_w(P, F)$  we have that  $|\partial P \cap N| \leq |\partial Q \cap N|$ .

Minimality is a local property of the mutation graph. It is certainly possible for there to exist more than one minimal polygon in a given mutation-equivalence class (see Example 4.4 below), however the number is finite. This is shown in Theorem 5.4 in the case when  $\mathcal{B} = \emptyset$ , and in Theorem 6.3 when  $\mathcal{B} \neq \emptyset$ . Given a polygon  $P \subset N_{\mathbb{Q}}$  one can easily construct a mutation-equivalent minimal polygon. Set  $P_0 := P$  and recursively define  $P_{i+1}$  as follows. Let  $\Gamma_i := \{P_i\} \cup \{\text{mut}_w(P_i, F) \mid \text{for all possible } w \in M\}$  where, as usual, we regard a polygon as being defined only up to  $GL_2(\mathbb{Z})$ -equivalence. Pick  $P_{i+1} \in \Gamma_i$  such that  $|\partial P_{i+1} \cap N| = \min\{|\partial Q \cap N| \mid Q \in \Gamma_i\}$ . If  $|\partial P_i \cap N| = |\partial P_{i+1} \cap N|$  we stop, and  $P_i$  is minimal. Notice that this process must terminate in a finite number of steps.

**Lemma 4.2** (Characterisation of minimality). *Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon. The following are equivalent:*

- (i)  $P$  is minimal;
- (ii)  $|\partial P \cap N| \leq |\partial Q \cap N|$  for every mutation  $Q := \text{mut}_w(P, F)$ ;
- (iii)  $|P^\circ \cap N| \leq |Q^\circ \cap N|$  for every mutation  $Q := \text{mut}_w(P, F)$ ;
- (iv)  $\text{Vol}(P) \leq \text{Vol}(Q)$  for every mutation  $Q := \text{mut}_w(P, F)$ ;
- (v)  $r_1 + \dots + r_n \leq s_1 + \dots + s_n$  for every mutation  $Q := \text{mut}_w(P, F)$ , where the  $r_i$  (respectively  $s_i$ ) are the Gorenstein indices of the primitive  $T$ -cones associated with  $P$  (respectively  $Q$ ).

*Proof.* Let  $C$  be an arbitrary cone (not necessarily a  $T$ -cone) corresponding to the cyclic quotient singularity  $\frac{1}{R}(a, b)$ . Let  $\rho_1, \rho_2 \in N$  be the primitive lattice vectors generating the rays of  $C$ . Recall that  $k = \gcd\{R, a + b\}$  is the length of the line segment  $\rho_1\rho_2$ , and that the Gorenstein index  $r = R/\gcd\{R, a + b\}$  is the height of  $\rho_1\rho_2$ . Set  $D := \text{conv}\{\rho_1, \rho_2, \mathbf{0}\}$ . Since  $R = k + 2|D^\circ \cap N|$  we have that:

$$|D^\circ \cap N| = \frac{R - k}{2} = \frac{k(r - 1)}{2}.$$

If  $C$  is a primitive  $T$ -cone then  $r = k$ .

Let  $P$  have singularity content  $(n, \mathcal{B})$ , and let  $r_1, \dots, r_n$  be the Gorenstein indices of the primitive  $T$ -cones. Then the number of boundary points is

$$(4.1) \quad |\partial P \cap N| = \sum_{i=1}^n r_i + \sum_{\mathcal{B}} (|\partial D \cap N| - 1),$$

where  $\sum_{\mathcal{B}}(|\partial D \cap N| - 1)$  is the contribution arising from the basket  $\mathcal{B}$ , and the number of interior points is given by

$$(4.2) \quad |P^\circ \cap N| = 1 + \frac{1}{2} \sum_{i=1}^n r_i(r_i - 1) + \sum_{\mathcal{B}} |D^\circ \cap N|,$$

where  $\sum_{\mathcal{B}} |D^\circ \cap N|$  is the contribution arising from the basket  $\mathcal{B}$ . Notice that the values of both  $\sum_{\mathcal{B}}(|\partial D \cap N| - 1)$  and  $\sum_{\mathcal{B}} |D^\circ \cap N|$  are fixed under mutation. By applying Pick's formula we obtain

$$(4.3) \quad \text{Vol}(P) = |\partial P \cap N| + 2|P^\circ \cap N| - 2 = \sum_{i=1}^n r_i^2 + B$$

where  $B := \sum_{\mathcal{B}}(|\partial D \cap N| - 1) + 2 \sum_{\mathcal{B}} |D^\circ \cap N|$  is a constant under mutation. Mutation can change the value of only one  $r_i$  at a time, hence equations (4.1), (4.2), and (4.3) are all locally minimal with respect to mutation if and only if  $r_1 + \dots + r_n$  is locally minimal with respect to mutation.  $\square$

**Remark 4.3.** Recall that mutations between Fano triangles are characterised in terms of solutions to a Diophantine equation [4]. Every solution can be obtained from a minimal solution – a solution  $(a, b, c) \in \mathbb{Z}_{>0}^3$  whose sum  $a + b + c$  is minimal – and a minimal solution corresponds to a triangle with smallest area [4, Lemma 3.16]. By Lemma 4.2 (iv) we see that the notion of minimality introduced in Definition 4.1 above can be viewed as a generalisation of the concept of minimal solution.

**Example 4.4.** Any reflexive polygon is minimal by Lemma 4.2 (iii), since  $|P^\circ \cap N| = 1$ . In particular this gives us examples of mutation-equivalent polygons  $P_1 \not\cong P_2$ , both of which are minimal. For example, one could take  $P_1 = \text{conv}\{(\pm 1, 0), (0, \pm 1)\}$  and  $P_2 = \text{conv}\{(1, 0), (0, 1), (-1, -2)\}$  (the polygons associated with  $\mathbb{P}^1 \times \mathbb{P}^1$  and  $\mathbb{P}(1, 1, 2)$ , respectively).

An immediate consequence of Lemma 4.2 (v) is the following:

**Corollary 4.5.** *Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon. For each edge  $E$  of  $P$  let  $w_E \in M$  denote the corresponding primitive inner normal vector and let  $k_E$  denote the length.  $P$  is minimal if and only if  $|h_{\min}| \leq h_{\max}$  for each  $w_E$  such that  $k_E \geq |h_{\min}|$ .*

**Example 4.6.** Any centrally symmetric polygon (that is, any polygon  $P$  satisfying  $v \in P$  if and only if  $-v \in P$ ) is minimal: for any primitive inner normal vector  $w \in M$ ,  $|h_{\min}| = h_{\max}$ .

**Corollary 4.7.** *Let  $P := \text{conv}\{v_0, v_1, v_2\}$  be a Fano triangle with residual basket  $\mathcal{B} = \emptyset$ . Then  $P$  is minimal if and only if  $v_0 + v_1 + v_2 \in P$ .*

*Proof.* Set  $\bar{v} := v_0 + v_1 + v_2 \in N$ . Let  $E$  be an edge of  $P$  and let  $w \in M$  be the corresponding primitive inner normal vector. Then  $w(\bar{v}) = w(v_0) + w(v_1) + w(v_2) = 2h_{\min} + h_{\max}$ , and  $|h_{\min}| \leq h_{\max}$  if and only if  $\bar{v}$  lies in the half-space  $H_E := \{v \in N_{\mathbb{Q}} \mid w(v) \geq h_{\min}\}$ . By Corollary 4.5 we have that  $P$  is minimal if and only if  $\bar{v} \in \bigcap H_E$ , where the intersection is taken over all edges  $E$  of  $P$ . The result follows.  $\square$

## 5. MINIMAL FANO POLYGONS WITH ONLY $T$ -SINGULARITIES

In Theorem 5.4 below we classify all minimal Fano polygons with residual basket  $\mathcal{B} = \emptyset$ . Conjecture A tells us that the mutation-equivalence classes should correspond to the ten qG-deformation classes of smooth del Pezzo surfaces, and in Theorem 5.6 we find that this is indeed what happens. In particular, in the case when  $X_P$  is a rank-one toric del Pezzo surface with only  $T$ -singularities we recover the results of Hacking–Prokhorov [25, Theorem 4.1]. Perling has also studied these surfaces from the viewpoint of mutations of exceptional collections [34].

**Definition 5.1.** Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon with vertex set denoted by  $\mathcal{V}(P)$  and edge set denoted by  $\mathcal{F}(P)$ . For each edge  $E \in \mathcal{F}(P)$  of  $P$  the unique primitive lattice point in the dual lattice  $M := \text{Hom}(N, \mathbb{Z})$  defining an inner normal of  $E$  is denoted by  $w_E$ . The positive integer  $r_E := -w_E(E)$  (that is, the height of  $E$  above the origin  $\mathbf{0}$ ) is equal to the Gorenstein index of the singularity associated with  $\text{cone}(E)$ . We call  $r_E$  the *Gorenstein index* (or *local index*) of  $E$ . The *maximum Gorenstein index* (also called the *maximum local index*) is the maximum Gorenstein index of all edges of  $P$ :

$$m_P := \max\{r_E \mid E \in \mathcal{F}(P)\}.$$

The Gorenstein index  $r_P$  of a Fano polygon  $P$  is the least common multiple of the Gorenstein indices:

$$r_P := \text{lcm}\{r_E \mid E \in \mathcal{F}(P)\}.$$

Equivalently,  $r_P$  is equal to the smallest positive integer  $\ell$  such that  $\ell P^*$  is a lattice polygon. In terms of the toric variety  $X_P$ ,  $r_P$  is equal to the smallest positive integer  $\ell$  such that  $-\ell K_{X_P}$  is Cartier.

**Lemma 5.2.** *Let  $P = \text{conv}\{v_0, v_1, v_2\} \subset N_{\mathbb{Q}}$  be a Fano triangle, and let  $(\lambda_0, \lambda_1, \lambda_2) \in \mathbb{Z}_{>0}^3$  be such that  $\lambda_0 v_0 + \lambda_1 v_1 + \lambda_2 v_2 = \mathbf{0}$ . Then:*

$$\text{Vol}(P) \cdot \text{Vol}(P^*) = \frac{(\lambda_0 + \lambda_1 + \lambda_2)^3}{\lambda_0 \lambda_1 \lambda_2}.$$

*Proof.* We give a toric proof; for a combinatorial argument, see [6, Proposition 6.2]. We can assume that the weights  $(\lambda_0, \lambda_1, \lambda_2)$  are coprime and, since  $P$  is a Fano polygon, this implies that the weights are pairwise coprime. Recall [27] that the Fano triangle  $P$  corresponds to some fake weighted projective space  $X_P = \mathbb{P}(\lambda_0, \lambda_1, \lambda_2)/G$ , where  $G = N/N'$  is the quotient of  $N$  by the sublattice  $N' = v_0 \cdot \mathbb{Z} + v_1 \cdot \mathbb{Z} + v_2 \cdot \mathbb{Z}$  generated by the vertices of  $P$ . The order  $|G|$ , or equivalently the index  $[N : N']$  of the sublattice  $N'$ , is called the *multiplicity* of  $P$ , and denoted by  $\text{mult}(P)$ . Let  $Q \subset N_{\mathbb{Q}}$  be the Fano triangle associated with weighted projective space  $X_Q = \mathbb{P}(\lambda_0, \lambda_1, \lambda_2)$ . Then

$$\text{Vol}(Q) = \lambda_0 + \lambda_1 + \lambda_2 \quad \text{and} \quad \text{Vol}(Q^*) = \frac{(\lambda_0 + \lambda_1 + \lambda_2)^2}{\lambda_0 \lambda_1 \lambda_2},$$

where the second value is simply the degree  $(-K_{X_Q})^2$  of the weighted projective space  $X_Q$ . But  $Q$  is  $GL_2(\mathbb{Z})$ -equivalent to the restriction of  $P$  to  $N'$ . Hence  $\text{Vol}(P) = \text{mult}(P) \cdot \text{Vol}(Q)$  and:

$$\text{Vol}(P^*) = \frac{1}{\text{mult}(P)} \cdot \text{Vol}(Q^*) = \frac{(\lambda_0 + \lambda_1 + \lambda_2)^3}{\lambda_0 \lambda_1 \lambda_2} \cdot \frac{1}{\text{Vol}(P)}. \quad \square$$

**Lemma 5.3** ([28, Lemma 2.1]). *Let  $P \subset N_{\mathbb{Q}}$  be a Fano polygon and let  $E$  be an edge with Gorenstein index equal to the maximum Gorenstein index of  $P$ ,  $r_E = m_P$ . Assume that there exists a non-vertex lattice point  $v \in \text{cone}(E)$  with  $w_E(v) = -1$ . For every lattice point  $v' \in P \cap N \setminus E$  we have that  $v + v' \in P \cap N$ . In particular, if  $E'$  is an edge of  $P$  that is not parallel to  $v$ , then  $|E' \cap N| \leq |E \cap N|$ .*

**Theorem 5.4.** *Let  $P \subset N_{\mathbb{Q}}$  be a minimal Fano polygon with residual basket  $\mathcal{B} = \emptyset$ . Then, up to  $GL_2(\mathbb{Z})$ -equivalence,  $P$  is one of the following 35 polygons:*

- (i) the five reflexive triangles  $R_1, \dots, R_5$  in Table 1;
- (ii) the eleven reflexive polygons  $R_6, \dots, R_{16}$  in Table 2;
- (iii) the nine non-reflexive triangles  $T_1, \dots, T_9$  in Table 1;
- (iv) the ten non-reflexive polygons  $P_1, \dots, P_{10}$  in Table 2.

*Proof.* We consider the cases when  $P$  is reflexive, that is  $m_P = 1$ , and when  $P$  is not reflexive, that is  $m_P > 1$ , separately.

$m_P = 1$ : Every reflexive polygon is minimal since  $|P^\circ \cap N| = 1$ . The classification of the reflexive polygons is well-known: up to  $GL_2(\mathbb{Z})$ -equivalence there are sixteen polygons [7, 37], of which five are triangles. This proves (i) and (ii).

$m_P > 1$ : Let  $E \in \mathcal{F}(P)$  be an edge of maximum Gorenstein index  $r_E = m_P > 1$ . By assumption the length of  $E$  is some multiple  $k \in \mathbb{Z}_{>0}$  of  $r_E$ , that is,  $|E \cap N| = kr_E + 1$ . Since the cone  $C = \text{cone}(E)$  is a union of  $k$  primitive  $T$ -cones, each of which contains exactly one point at height  $-1$  with respect to  $w_E$ , there exist  $k$  distinct points  $v_1, \dots, v_k \in C^\circ \cap N$  such that  $w_E(v_i) = -1$ . After suitable change of basis we can insist that  $E = \text{conv}\{(-a, r_E), (b, r_E)\}$  where  $a, b > 0$ ,  $a < r_E$ , and  $a + b = kr_E$ . Hence the point  $((i-1)r_E, r_E)$  lies in the strict interior of  $E$  and so, after possible reordering,  $v_i = (i-1, 1)$ , for each  $i = 1, \dots, k$ .

Let  $v \in \mathcal{V}(P)$  be a vertex of  $P$  such that  $w_E(v) = \max\{w_E(v') \mid v' \in P\} = h_{\max}$ . Since  $P$  is minimal by assumption, we have that  $w_E(v) \geq r_E = |h_{\min}|$  (Corollary 4.5). By Lemma 5.3 we have that  $v$  is contained in the strip  $E - v_1 \cdot \mathbb{Z}_{>0}$ . Hence we can write  $v = (\alpha, -\beta)$  for some  $\beta \geq r_E$  and  $-a \leq \alpha \leq b$ . If  $k > 1$  then we have  $v \in E - v_k \cdot \mathbb{Z}_{>0}$  and so

$$(5.1) \quad -a \leq \alpha \leq b - (k-1)(r_E + \beta) \leq b - 2(k-1)r_E = -a - (k-2)r_E.$$

TABLE 1. The minimal Fano triangles  $P \subset N_{\mathbb{Q}}$  with singularity content  $(n, \emptyset)$ , vertices  $\mathcal{V}(P)$ , maximum Gorenstein index  $m_P$ , and weights  $(\lambda_0, \lambda_1, \lambda_2)$ , up to the action of  $GL_2(\mathbb{Z})$ . The corresponding toric varieties  $X_P$  and degrees  $(-K_{X_P})^2 = 12 - n$  are also given. Those  $P$  marked with  $\star$  are chosen as the representative polygon in the mutation-equivalence class, and are depicted in Figure 1 (page 19).

$P$	$\mathcal{V}(P)$	$\lambda_0$	$\lambda_1$	$\lambda_2$	$X_P$	$m_P$	$n$	$(-K_{X_P})^2$
$\star R_1$	$(1, 0), (0, 1), (-1, -1)$	1	1	1	$\mathbb{P}^2$	1	3	9
$\star R_2$	$(1, 1), (-2, 1), (1, -2)$	1	1	1	$\mathbb{P}^2/(\mathbb{Z}/3)$	1	9	3
$T_1$	$(1, 3), (-2, 3), (1, -6)$	1	1	1	$\mathbb{P}^2/(\mathbb{Z}/9)$	3	11	1
$R_3$	$(1, 1), (-1, 1), (0, -1)$	1	1	2	$\mathbb{P}(1, 1, 2)$	1	4	8
$R_4$	$(1, 1), (-1, 1), (1, -3)$	1	1	2	$\mathbb{P}(1, 1, 2)/(\mathbb{Z}/2)$	1	8	4
$T_2$	$(1, 2), (-1, 2), (1, -6)$	1	1	2	$\mathbb{P}(1, 1, 2)/(\mathbb{Z}/4)$	2	10	2
$\star T_3$	$(3, 2), (-1, 2), (-1, -2)$	1	1	2	$\mathbb{P}(1, 1, 2)/(\mathbb{Z}/4)$	2	10	2
$T_4$	$(1, 4), (-3, 4), (1, -4)$	1	1	2	$\mathbb{P}(1, 1, 2)/(\mathbb{Z}/8)$	4	11	1
$R_5$	$(1, 1), (-1, 1), (1, -2)$	1	2	3	$\mathbb{P}(1, 2, 3)$	1	6	6
$T_5$	$(1, 2), (-1, 2), (1, -4)$	1	2	3	$\mathbb{P}(1, 2, 3)/(\mathbb{Z}/2)$	2	9	3
$T_6$	$(1, 3), (-2, 3), (1, -3)$	1	2	3	$\mathbb{P}(1, 2, 3)/(\mathbb{Z}/3)$	3	10	2
$\star T_7$	$(5, 3), (-1, 3), (-1, -3)$	1	2	3	$\mathbb{P}(1, 2, 3)/(\mathbb{Z}/6)$	3	11	1
$T_8$	$(1, 2), (-1, 2), (1, -3)$	1	4	5	$\mathbb{P}(1, 4, 5)$	2	7	5
$T_9$	$(2, 5), (-3, 5), (2, -5)$	1	4	5	$\mathbb{P}(1, 4, 5)/(\mathbb{Z}/5)$	5	11	1

We conclude that  $k \leq 2$ .

$k = 2$ : Let us first consider the case when  $k = 2$ . By (5.1)  $P$  is a triangle given by

$$P = \text{conv}\{(-a, r_E), (-a + 2r_E, r_E), (-a, -r_E)\}, \quad \text{where } 0 < a < r_E, \gcd\{a, r_E\} = 1.$$

Let  $E' \in \mathcal{F}(P)$  be the edge with vertices  $(-a, r_E)$  and  $(-a, -r_E)$ . Since the corresponding cone is of class  $T$ , we have that  $a \mid 2r_E$ , and so  $a = 1$  or  $2$ . Analogously, by considering the edge with vertices  $(-a, -r_E)$  and  $(2r_E - a, r_E)$  of length  $2r_E$  and Gorenstein index  $r_E - a$ , we see  $r_E - a \mid 2r_E$ . This leads to  $(r_E, a) \in \{(2, 1), (3, 1), (3, 2), (4, 2), (6, 2)\}$ . The first possibility gives the triangle  $T_3$  in Table 1, the second and third possibilities define the triangle  $T_7$ , and the final two possibilities are excluded because of  $\gcd\{a, r_E\} = 1$ .

$k = 1$ : We now consider the case when  $k = 1$ . This is subdivided into two cases depending on whether or not there exists an edge  $E' \in \mathcal{F}(P)$  parallel to  $E$ .

Edge  $E'$  parallel to  $E$ : Let us assume that there exists a second point of  $P$  at height  $w_E(v)$ ; that is, that there exists an edge  $E' \in \mathcal{F}(P)$  such that  $w_E(E') = w_E(v)$ , so that  $E$  and  $E'$  are parallel. By minimality we see that  $r_{E'} = r_E$ , and by Lemma 5.3 we have that  $|E' \cap N| \leq |E \cap N| = r_E + 1$ . Recalling that cone( $E'$ ) is a  $T$ -cone, and hence  $r_{E'} \mid |E' \cap N| - 1$ , we see that  $P$  is a rectangle:

$$P = \text{conv}\{(-a, r_E), (-a + r_E, r_E), (-a, -r_E), (-a + r_E, -r_E)\}.$$

Since  $P$  is minimal we have that  $2a = r_E$  and, by primitivity of the vertices,  $a = 1, r_E = 2$ , giving  $P_1$  in Table 2.

No edge parallel to  $E$ : We are now in the situation where  $v$  is the unique vertex satisfying  $w_E(v) = \max\{w_E(v') \mid v' \in P\}$ . We subdivide this into two cases depending on whether one of the edges  $E' \in \mathcal{F}(P)$  with vertex  $v$  is vertical.

$E'$  not vertical: Let us assume that there is no vertical edge adjacent to  $v = (\alpha, -\beta)$ . We consider the case  $\alpha \geq 0$  (the case  $\alpha < 0$  being similar). By our assumption, we can choose an edge  $E'$  adjacent to  $v$  with  $w_{E'} = (-\gamma, \delta) \in M$ , where  $\gamma, \delta \geq 1$ . We have that  $r_E \geq -w_{E'}(v) = \gamma\alpha + \delta\beta \geq \alpha + \beta \geq \beta \geq r_E$ .



TABLE 2. The minimal Fano  $m$ -gons,  $m \geq 4$ ,  $P \subset N_{\mathbb{Q}}$  with singularity content  $(n, \emptyset)$ , vertices  $\mathcal{V}(P)$ , and maximum Gorenstein index  $m_P$ , up to the action of  $GL_2(\mathbb{Z})$ . The degrees  $(-K_{X_P})^2 = 12 - n$  of the corresponding toric varieties are also given. Those  $P$  marked with  $\star$  are chosen as the representative polygon in the mutation-equivalence class, and are depicted in Figure 1 (page 19).

$P$	$\mathcal{V}(P)$	$m_P$	$n$	$(-K_{X_P})^2$
$P_1$	$(1, 2), (-1, 2), (-1, -2), (1, -2)$	2	10	2
$P_2$	$(1, 2), (-1, 2), (-1, -1), (1, -3)$	2	10	2
$P_3$	$(1, 2), (-1, 2), (-1, 0), (1, -4)$	2	10	2
$P_4$	$(1, 2), (-1, 2), (-1, 1), (1, -5)$	2	10	2
$P_5$	$(1, 2), (-1, 2), (-1, 0), (1, -2)$	2	9	3
$P_6$	$(1, 2), (-1, 2), (-1, 1), (1, -3)$	2	9	3
$R_6$	$(1, 1), (-1, 1), (-1, 0), (1, -2)$	1	8	4
$\star R_7$	$(1, 1), (-1, 1), (-1, -1), (1, -1)$	1	8	4
$P_7$	$(1, 2), (-1, 2), (0, -1), (1, -3)$	2	8	4
$P_8$	$(1, 2), (-1, 2), (-1, 1), (0, -1), (1, -2)$	2	8	4
$R_8$	$(1, 1), (-1, 1), (0, -1), (1, -2)$	1	7	5
$P_9$	$(1, 2), (-1, 2), (-1, 1), (1, -2)$	2	7	5
$P_{10}$	$(1, 2), (-1, 2), (0, -1), (1, -2)$	2	7	5
$\star R_9$	$(1, 1), (-1, 1), (-1, 0), (0, -1), (1, -1)$	1	7	5
$R_{10}$	$(1, 1), (-1, 1), (-1, 0), (1, -1)$	1	6	6
$R_{11}$	$(1, 0), (1, 1), (-1, 1), (-1, 0), (0, -1)$	1	6	6
$\star R_{12}$	$(1, 0), (1, 1), (0, 1), (-1, 0), (-1, -1), (0, -1)$	1	6	6
$R_{13}$	$(1, 0), (1, 1), (-1, 1), (0, -1)$	1	5	7
$\star R_{14}$	$(1, 0), (1, 1), (0, 1), (-1, -1), (0, -1)$	1	5	7
$\star R_{15}$	$(1, 0), (0, 1), (-1, -1), (0, -1)$	1	4	8
$\star R_{16}$	$(1, 0), (0, 1), (-1, 0), (0, -1)$	1	4	8

Hence  $(\alpha, -\beta) = (0, -r_E)$ ,  $w_{E'} = (-\gamma, 1)$ , and  $r_{E'} = r_E = m_P$ . Exchanging the roles of  $E$  and  $E'$  we have that  $E'$  is of length either  $2r_E$ , in which case we are in the case  $k = 2$  above, or of length  $r_E$ , in which case  $(r_E, (\gamma - 1)r_E)$  is a vertex of  $P$ . This is a contradiction, since as  $r_E > b$ , this vertex is not contained in the strip  $E - v_1 \cdot \mathbb{Z}_{>0}$ .

$E'$  vertical: The majority of cases arise when  $v$  is contained in a vertical edge  $E' \in \mathcal{F}(P)$ . This edge necessarily contains one of the two vertices of  $E$ , and without loss of generality (since  $a + b = r_E$ ) we may assume that  $E' = \text{conv}\{v, (b, r_E)\}$ . Hence  $v = (b, r_E - jb)$  for some  $j \in \mathbb{Z}_{>0}$ . Minimality forces  $r_E - jb \leq -r_E$ , so that

$$(5.2) \quad jb \geq 2r_E.$$

Moreover, minimality implies

$$(5.3) \quad 2b \leq r_E.$$

In particular we see that  $j \geq 4$ .

$j = 4$ : The case when  $j = 4$  is different from the cases when  $j \geq 5$ , and we deal with it now. Equations (5.2) and (5.3) gives that  $2b = r_E$ , and so by primitivity we have that  $r_E = 2$  and  $b = 1$ .

Hence  $P$  is contained in the rectangle  $[-1, 1] \times [-2, 2]$ . Notice that the requirement that  $\mathbf{0} \in P^\circ$  means that  $P$  cannot be a triangle. We find  $P_5, P_8, P_9$ , and  $P_{10}$  in Table 2.

$j \geq 5$ : Consider the triangle

$$T := \text{conv}\{(b - r_E, r_E), (b, r_E), (b, r_E - jb)\}.$$

By (5.2)  $-r_E \geq r_E - jb$ , so the point  $(b, -r_E)$  is in  $T$ . The midpoint of  $(b, -r_E)$  and  $(b - r_E, r_E)$  is  $(b - r_E/2, 0)$ . Equation (5.3) implies  $b - r_E/2 \leq 0$ . If  $b - r_E/2 < 0$  then  $\mathbf{0} \in T^\circ$ . If  $b - r_E/2 = 0$  then  $j \geq 5$  implies that  $(b, -r_E)$  is not equal to the vertex  $(b, r_E - jb)$ , so  $\mathbf{0} = (b - r_E/2, 0) \in T^\circ$  again.

This shows that  $T$  is a Fano triangle. We have that  $\text{Vol}(T) = jbr_E$ . Moreover, one can check that

$$(jb^2, jbr_E - jb^2 - r_E^2, r_E^2) \in \mathbb{Z}_{>0}^3$$

satisfy the conditions of Lemma 5.2. Hence

$$\text{Vol}(T^*) = \frac{j}{jbr_E - jb^2 - r_E^2}.$$

By Proposition 3.8 (note  $\text{Vol}(P^*) = (-K_{X_P})^2$  and  $\mathcal{B} = \emptyset$ ),  $\text{Vol}(P^*) \in \mathbb{Z}_{>0}$ , and so  $1 \leq \text{Vol}(P^*) \leq \text{Vol}(T^*)$ . Let  $b = r_E q$ , where  $\frac{2}{j} \leq q \leq \frac{1}{2}$ , by (5.2) and (5.3), so that the lower bound on  $\text{Vol}(T^*)$  gives:

$$(5.4) \quad r_E^2(-jq^2 + jq - 1) \leq j.$$

The quadratic in  $q$  on the left-hand-side of (5.4) is strictly positive in the range  $\frac{2}{j} \leq q \leq \frac{1}{2}$  and obtains its minimum value when  $q = \frac{2}{j}$ . Hence (5.4) gives us:

$$(5.5) \quad r_E^2 \leq \frac{j^2}{j - 4}.$$

Recall from Proposition 3.8 that  $\text{Vol}(P^*) = 12 - n \geq 1$ , where  $n$  is the total number of primitive  $T$ -cones spanned by the edges of  $P$ ; equivalently,

$$n = \sum_{F \in \mathcal{F}(P)} \frac{|F \cap N| - 1}{r_F}.$$

Since  $P$  must have at least three edges, each of which corresponds to a  $T$ -cone, and by construction we have that the top edge decomposes into a single primitive  $T$ -cone, and that the right-hand vertical edge  $E'$  decomposes into  $j \geq 5$  primitive  $T$ -cones, we conclude that  $j \in \{5, \dots, 9\}$ . From the inequalities (5.2), (5.3), (5.5), and  $m_P = r_E \geq 2$ , along with the requirement that  $\text{gcd}\{b, r_E\} = 1$ , we obtain finitely many possibilities for the triple  $(j, r_E, b)$ , as recorded in Table 3.

TABLE 3. The possible values of  $(j, r_E, b)$ .


$j$	5	5	6	6	7	7	8	8	8	9	9	9
$r_E$	2	5	2	3	2	3	2	3	4	2	3	4
$b$	1	2	1	1	1	1	1	1	1	1	1	1

Analysis of Table 3: First we consider the cases where  $r_E = 2$  and  $b = 1$ . Here  $P$  is contained in the rectangle  $[-1, 1] \times [-j + 2, 2]$ . Let  $E''$  be the edge with  $(0, -1) \in \text{cone}(E'')$  that contains the lower right vertex  $(1, 2 - j) \in E''$ . If  $(0, -1) \notin P$  then one immediately finds that  $j = 5$  and  $P$  is a triangle. This is the triangle  $T_8$  in Table 1. If  $(0, -1) \in \partial P$  then  $j \leq 6$ , and  $P$  is either the triangle  $T_5$ , or one of  $P_6$  or  $P_7$  in Table 2. Finally, if  $(0, -1) \in P^\circ$  then the Gorenstein index of  $E''$  must be two. Therefore  $(0, -2) \in E''$ , giving either the triangle  $T_2$ , or one of  $P_2, P_3$ , or  $P_4$ .

In the cases  $(5, 5, 2)$ ,  $(8, 4, 1)$ , and  $(9, 3, 1)$ , we have equality in (5.4), hence  $1 = \text{Vol}(P^*) = \text{Vol}(T^*)$  and so  $P = T$  is uniquely determined. This gives the triangles  $T_1, T_4$ , and  $T_9$ . In the case  $(6, 3, 1)$  the triangle  $T = T_6$ . In the remaining cases it is easily verified that  $T$  is not a minimal triangle with only  $T$ -singularities; this completes the proof of (iii).

Consider the cases  $(6, 3, 1)$ ,  $(7, 3, 1)$ , and  $(8, 3, 1)$ . In all three cases  $r_E = 3$  and  $b = 1$ , hence  $P$  is contained in the rectangle  $[-2, 1] \times [-j + 3, 3]$ ,  $j \in \{6, 7, 8\}$ . If we assume that  $P$  is not a triangle,

FIGURE 1. A representative minimal Fano polygon  $P \subset N_{\mathbb{Q}}$  with singularity content  $(n, \emptyset)$ , for each of the 10 mutation-equivalence classes. The representatives correspond, up to the action of  $GL_2(\mathbb{Z})$ , with the polygons marked  $\star$  in Tables 2 and 4. The corresponding toric varieties have degree  $(-K_{X_P})^2 = 12 - n$ .



it follows that that  $(0, -1) \in P^\circ$ , therefore  $(0, -1) \in \text{cone}(E'')$  for an edge  $E'' \in \mathcal{F}(P)$  containing the bottom-right vertex  $(1, 3 - j)$ . This implies that the edge  $E''$  has Gorenstein index two or three. Let us assume the Gorenstein index of  $E''$  is three. Then it must have length three, and so  $P$  is not a triangle; there must exist one more vertex with first coordinate  $-2$ . However, this means that there exists a left vertical edge, contradicting minimality. Now assume that the Gorenstein index of  $E''$  is two. Hence  $(0, -2)$  is an interior point in  $E''$ , and since  $E''$  has length two there is a unique vertex with first coordinate  $-1$ . Hence there can be at most one vertex left with first coordinate  $-2$ , excluding the vertex  $(-2, 3)$ . Enumerating these possibilities shows that none result in a minimal Fano polygon with only  $T$ -cones.


Finally, consider the case  $(9, 4, 1)$ . We see that  $(0, -2) \in P^\circ$ , hence the non-vertical edge  $E'' \in \mathcal{F}(P)$  containing  $(1, -5)$  is of Gorenstein index either three or four. If  $r_{E''} = 3$  then  $(-2, 1)$  is a vertex of  $P$ , and if  $r_{E''} = 4$  then  $(-3, -1)$  is a vertex of  $P$ . In either case  $P$  fails to have only  $T$ -cones, and so (iv) is complete.  $\square$

**Corollary 5.5.** *There are precisely two mutation-equivalence classes of Fano polygons with singularity content  $(4, \emptyset)$ . These classes are given by  $R_{15}$  and  $R_{16}$ , corresponding to the toric varieties  $\mathbb{F}_1$  and  $\mathbb{P}^1 \times \mathbb{P}^1$ , respectively.*

*Proof.* By Theorem 5.4 there are only three minimal polygons with singularity content  $(4, \emptyset)$ :  $R_3$ ,  $R_{15}$ , and  $R_{16}$ . It is easy to see that  $R_3$  and  $R_{16}$  are mutation-equivalent. That  $R_{15}$  and  $R_{16}$  are distinct up to mutation is shown in Example 3.12.  $\square$

**Theorem 5.6.** *There are ten mutation-equivalence classes of Fano polygon with residual basket  $\mathcal{B} = \emptyset$ . Representative polygons for each mutation-equivalence class are given by  $R_1, R_{16}, R_{15}, R_{14}, R_{12}, R_9, R_7, R_2, T_3$ , and  $T_7$ . These representatives are depicted in Figure 1.*

*Proof.* Let  $P, Q \subset N_{\mathbb{Q}}$  be two minimal Fano polygons as given in Theorem 5.4 with  $\text{SC}(P) = \text{SC}(Q)$ . With the exception of the case when  $\text{SC}(P) = \text{SC}(Q) = (4, \emptyset)$ , which is handled in Corollary 5.5 above, it can easily be seen that  $P$  and  $Q$  are mutation-equivalent. We will do the case when  $\text{SC}(P) = \text{SC}(Q) = (6, \emptyset)$ ; the remaining cases are similar. The minimal Fano polygons  $R_5, R_{10}, R_{11}$ , and  $R_{12}$  are connected via a sequence of mutations:



The mutations have been labelled with their corresponding primitive inner normal vector  $w$ .  $\square$

## 6. FINITENESS OF MINIMAL FANO POLYGONS

In this section we generalise Theorem 5.4 to the case when the residual basket  $\mathcal{B} \neq \emptyset$ .

**Definition 6.1.** Given a residual basket  $\mathcal{B} \neq \emptyset$  we define

$$\begin{aligned} m_{\mathcal{B}} &:= \max\{r_{\sigma} \mid \sigma \in \mathcal{B}\}, \\ d_{\mathcal{B}} &:= \text{lcm}\{\text{denom}(A_{\sigma}) \mid \sigma \in \mathcal{B}\}, \\ s_{\mathcal{B}} &:= -\min(\{0\} \cup \{A_{\sigma} \mid \sigma \in \mathcal{B}\}), \end{aligned}$$

where  $r_{\sigma}$  is the Gorenstein index of  $\sigma$ ,  $A_{\sigma}$  is the contribution of  $\sigma$  to the degree (as given in Proposition 3.8), and  $\text{denom}(x)$  denotes the denominator of  $x \in \mathbb{Q}$ . In the case when  $\mathcal{B} = \emptyset$  we define  $m_{\mathcal{B}} := 1$ ,  $d_{\mathcal{B}} := 1$ , and  $s_{\mathcal{B}} := 0$ .

**Remark 6.2.** Bounding  $m_{\mathcal{B}}$  automatically bounds the number of possible types of singularities that can occur in the residual basket. In particular there are only finitely many possible values of  $A_{\sigma}$ , hence  $d_{\mathcal{B}}$  and  $s_{\mathcal{B}}$  are bounded from above.

**Theorem 6.3.** *There exist only a finite number of minimal Fano polygons, up to the action of  $GL_2(\mathbb{Z})$ , with bounded maximum Gorenstein index  $m_{\mathcal{B}}$  of the cones in the residual baskets  $\mathcal{B}$ .*

*Proof.* We assume throughout that  $\mathcal{B} \neq \emptyset$ , the empty case having already been considered in Theorem 5.4. The proof is constructive, and follows a similar structure to the proof when  $\mathcal{B} = \emptyset$ .

$m_P = m_{\mathcal{B}}$ : The number of possible Fano polygons with bounded maximum Gorenstein index  $m_P$  is known to be finite, up to  $GL_2(\mathbb{Z})$ -equivalence. Algorithms for computing all such Fano polygons are described in [28].

$m_P > m_{\mathcal{B}}$ : Let  $E \in \mathcal{F}(P)$  be an edge of maximum Gorenstein index  $r_E = m_P > m_{\mathcal{B}}$ . In particular  $\text{cone}(E)$  is a  $T$ -cone, hence  $|E \cap N| = kr_E + 1$  for some  $k \in \mathbb{Z}_{>0}$ . After suitable change of basis we can insist that  $E = \text{conv}\{(-a, r_E), (b, r_E)\}$  where  $a, b > 0$ ,  $a < r_E$ , and  $a + b = kr_E$ . Hence, as in the proof of Theorem 5.4, there exist  $k$  distinct points  $v_i = (i - 1, 1)$ , for  $i = 1, \dots, k$ , where each  $v_i \in \text{cone}(E)^\circ$ ,  $w_E(v_i) = -1$ . Let  $v \in \mathcal{V}(P)$  be a vertex of  $P$  such that  $w_E(v) = \max\{w_E(v') \mid v' \in P\}$ . Since  $P$  is minimal by assumption, by Corollary 4.5 we have that  $w_E(v) \geq r_E$ . By applying Lemma 5.3 with respect to  $v_1$  and  $v_k$  we conclude that  $k \leq 2$ .

$k = 2$ : First we consider the case when  $k = 2$ . As in the proof of Theorem 5.4 we have that

$$P = \text{conv}\{(-a, r_E), (-a + 2r_E, r_E), (-a, -r_E)\}, \quad \text{where } 0 < a < r_E, \text{gcd}\{a, r_E\} = 1.$$

Let  $E_1$  and  $E_2$  be the two edges of  $P$  distinct from the horizontal edge  $E$ , where  $r_{E_1} = a$  and  $r_{E_2} = r_E - a$ . Since at least one of these two edges has Gorenstein index  $m_{\mathcal{B}}$ , by symmetry we can assume that  $a = m_{\mathcal{B}}$ . The edge  $E_2$  is of length  $2r_E$ , giving  $2r_E = j(r_E - m_{\mathcal{B}}) + l$  for some  $j \in \mathbb{Z}_{\geq 0}$ ,  $0 \leq l < r_E - m_{\mathcal{B}}$ .

If  $l = 0$  then  $jm_{\mathcal{B}} = (j - 2)r_E$ , and  $\text{gcd}\{m_{\mathcal{B}}, r_E\} = 1$  implies that  $r_E \mid j$ . Writing  $j = j'r_E$  for some  $j' \in \mathbb{Z}_{\geq 0}$  we see that  $2 = j'(r_E - m_{\mathcal{B}})$  and hence  $r_E = m_{\mathcal{B}} + 1$  or  $r_E = m_{\mathcal{B}} + 2$ . If  $l > 0$  then  $r_E - m_{\mathcal{B}} \leq m_{\mathcal{B}}$ , and so  $r_E < 2m_{\mathcal{B}}$  (the case of equality being excluded by primitivity). Hence in either case the number of possible minimal polygons is finite.

$k = 1$ : We now consider the case when  $k = 1$ . Once again we subdivide this into two cases depending on whether there exists an edge  $E' \in \mathcal{F}(P)$  parallel to  $E$ .

Edge  $E'$  parallel to  $E$ : First we assume that there exists an edge  $E' \in \mathcal{F}(P)$  such that  $v$  is a vertex of  $\overline{E'}$ , and  $E'$  is parallel to  $E$ . If  $\text{cone}(E')$  contains a residual component then  $r_{E'} \leq m_{\mathcal{B}}$ . By minimality  $r_E \leq m_{\mathcal{B}}$ , which is a contradiction. Therefore  $\text{cone}(E')$  must be of class  $T$ , and by minimality we see that  $r_{E'} = r_E$ . As in the proof of Theorem 5.4 we conclude that  $P$  is a rectangle:

$$P = \text{conv}\{(-a, r_E), (-a + r_E, r_E), (-a, -r_E), (-a + r_E, -r_E)\}.$$

Since one of the two vertical edges must contain a residual component at height  $m_{\mathcal{B}}$ , without loss of generality we may assume that  $a = m_{\mathcal{B}}$ , hence  $2r_E = jm_{\mathcal{B}} + l$  for some  $j \in \mathbb{Z}_{\geq 0}$ ,  $0 < l < m_{\mathcal{B}}$ . Notice that if  $j = 0$  then  $2r_E = l < m_{\mathcal{B}}$ , a contradiction. Hence  $j > 0$ . The second vertical edge lies at height

$r_E - m_B$ , and by Corollary 4.5 we have that  $2m_B \leq r_E$ . Hence  $2r_E = j'(r_E - m_B) + l'$  for some  $j' \in \mathbb{Z}_{\geq 0}$ ,  $0 \leq l' < r_E - m_B$ . If  $j' = 0$  then  $2r_E = l' < r_E - m_B$ , a contradiction. If  $j' > 0$  then, by minimality,  $r_E \leq 2m_B$ , implying that  $r_E = 2m_B$ . But this contradicts primitivity, hence this case does not occur.

No edge parallel to  $E$ : We now assume that  $v$  is the unique point in  $P$  such that  $w_E(v) = \max\{w_E(v') \mid v' \in P\}$ . Once again we subdivide this into two cases, depending on whether there exists a vertical edge  $E' \in \mathcal{F}(P)$  with vertex  $v$ .

$E'$  not vertical: This proof in this case is identical to that of Theorem 5.4: it results in no minimal polygons.

$E'$  vertical: Without loss of generality we may assume that there is a vertical edge  $E' \in \mathcal{F}(P)$  with vertices  $v$  and  $(b, r_E)$ . Hence  $v = (b, r_E - jb - l)$  for some  $j \in \mathbb{Z}_{\geq 0}$ ,  $0 \leq l < b$ . By minimality of  $E$ ,

$$(6.1) \quad 2r_E \leq jb + l.$$

Notice that if  $j = 0$  then  $2r_E \leq l < m_B$ , a contradiction. Hence  $j > 0$  and, by minimality of  $E'$ ,

$$(6.2) \quad 2b \leq r_E.$$

$l = 0$ : Consider the case when  $l = 0$ . Inequalities (6.1) and (6.2) imply that  $j \geq 4$ .

$j = 4$ : Assume that  $j = 4$ . Then  $2b = r_E$ , so  $\gcd\{b, r_E\} = 1$  implies that  $r_E = 2$  and  $b = 1$ . Hence  $P$  is contained in the rectangle  $[-1, 1] \times [-2, 2]$ . This contains only four possible polygons, all of which have  $m_P \leq 3$ , contradicting  $m_P > m_B \geq 3$ . Hence this case does not occur.

$j \geq 5$ : As in the proof of Theorem 5.4,  $\mathbf{0}$  is contained in the strict interior of the triangle

$$T := \text{conv}\{(b - r_E, r_E), (b, r_E), (b, r_E - jb)\} \subset P.$$

As before, the area of the dual triangle is given by

$$\text{Vol}(T^*) = \frac{j}{jbr_E - jb^2 - r_E^2}.$$

By Proposition 3.8,  $\text{Vol}(P^*) \in \frac{1}{d_B} \cdot \mathbb{Z}_{>0}$ , hence  $\frac{1}{d_B} \leq \text{Vol}(P^*) \leq \text{Vol}(T^*)$ . Let  $b = r_E q$ , where  $\frac{2}{j} \leq q \leq \frac{1}{2}$ , by (6.1) and (6.2). Then  $r_E^2(-jq^2 + jq - 1) \leq jd_B$ , and by considering the minimum value achieved by the quadratic in  $q$  on the left-hand-side of this inequality we obtain

$$(6.3) \quad r_E^2 \leq \frac{j^2 d_B}{j - 4} = \left( j + 4 + \frac{16}{j - 4} \right) d_B \leq (j + 20) d_B.$$

Notice that the number of edges of  $P$  distinct from  $E$  and  $E'$  can be at most  $r_E + 1$ . By Proposition 3.8 we obtain that  $12 - 1 - j + (r_E + 1)s_B \geq \text{Vol}(P^*) > 0$ . Hence

$$(6.4) \quad j < 11 + (r_E + 1)s_B.$$

Combining inequalities (6.3) and (6.4) gives  $r_E^2 < (31 + (r_E + 1)s_B)d_B$ , which implies that

$$(6.5) \quad r_E < \frac{s_B d_B + \sqrt{(s_B d_B)^2 + 4d_B(s_B + 31)}}{2}.$$

Since  $j$  is bounded by (6.4), the result follows.

$l > 0$ : Finally, let us consider the case when the edge  $E'$  contributes a residual singularity to the basket  $\mathcal{B}$ . Notice that if we had equality in (6.2), primitivity forces  $r_E = 2$  and  $b = 1$ , and so  $l = 0$ , which is a contradiction. Hence we conclude that the inequality is strict:

$$(6.6) \quad 2b < r_E.$$

Once again we find that  $j \geq 4$ .

Consider the triangle

$$T := \text{conv}\{(b - r_E, r_E), (b, r_E), (b, r_E - jb - l)\} \subset P.$$

One can check that  $\mathbf{0} \in T^\circ$ , so  $T$  is a Fano triangle with  $\text{Vol}(T) = r_E(jb + l)$ . Moreover, the weights

$$(b(jb + l), (jb + l)(r_E - b) - r_E^2, r_E^2) \in \mathbb{Z}_{>0}^3$$

satisfy the conditions of Lemma 5.2 (that the second weight is strictly positive follows from (6.1) and (6.6):  $(jb + l)(r_E - b) - r_E^2 \geq 2r_E^2 - 2r_E b - r_E^2 = r_E(r_E - 2b) > 0$ ). Hence

$$\text{Vol}(T^*) = \frac{jb + l}{b(jb + l)(r_E - b) - br_E^2} < \frac{j + 1}{(jb + l)(r_E - b) - r_E^2}.$$

Recalling, as above, that  $\frac{1}{d_B} \leq \text{Vol}(P^*) \leq \text{Vol}(T^*)$  we obtain:

$$(6.7) \quad (jb + l)(r_E - b) - r_E^2 < (j + 1)d_B.$$

The quadratic in  $b$  on the left-hand-side of (6.7) is strictly positive in the range  $\frac{2r_E - l}{j} \leq b < \frac{r_E}{2}$ , and obtains its minimum value when  $b = \frac{2r_E - l}{j}$ . Hence (6.7) gives:

$$(6.8) \quad (j - 4)r_E^2 + 2lr_E < j(j + 1)d_B.$$

We consider the cases  $j = 4$  and  $j > 4$  separately.

$j = 4$ : When  $j = 4$  inequalities (6.1) and (6.6) give  $r_E - \frac{1}{2} \leq 2b < r_E$ . Hence if  $l = 1$  this case does not occur. If  $l > 1$  then (6.8) gives us that  $r_E < 10d_B$ , resulting in only finitely many possibilities.

$j \geq 5$ : When  $j \geq 5$  inequality (6.8) implies that

$$(6.9) \quad r_E^2 < \frac{j(j + 1)d_B}{j - 4} = \left(j + 5 + \frac{20}{j - 4}\right) d_B \leq (j + 25)d_B.$$

Notice that the number of edges of  $P$  distinct from  $E$  and  $E'$  is at most  $r_E + 1$ , so by Proposition 3.8 we have that  $12 - 1 - j + (r_E + 2)s_B \geq \text{Vol}(P^*) > 0$ . Hence

$$(6.10) \quad j < 11 + (r_E + 2)s_B.$$

Combining inequalities (6.9) and (6.10) gives  $r_E^2 < (36 + (r_E + 2)s_B)d_B$ . This implies that

$$(6.11) \quad r_E < \frac{s_B d_B + \sqrt{(s_B d_B)^2 + 4d_B(2s_B + 36)}}{2}.$$

Since  $j$  is bounded by (6.10) we have only finitely many possible minimal polygons.  $\square$

## 7. MINIMAL FANO POLYGONS WITH $\mathcal{B} = \{m \times \frac{1}{3}(1, 1)\}$

In this section we apply Theorem 6.3 in order to classify all minimal Fano polygons with residual basket  $\mathcal{B}$  containing only singularities of type  $\frac{1}{3}(1, 1)$ . We find 65 minimal Fano polygons (Theorem 7.1 and Table 4), which result in 26 mutation-equivalence classes (Theorem 7.4 and Figure 2). These mutation-equivalence classes correspond exactly with the classification of Corti–Heuberger of qG-deformation-equivalence classes of del Pezzo surfaces of class TG with  $m \times \frac{1}{3}(1, 1)$  singular points [14].

In some sense  $\frac{1}{3}(1, 1)$  is the “simplest” residual singularity. Up to change of basis, the corresponding cone is given by  $C := \text{cone}\{(1, 0), (2, 3)\}$ . The length of the line segment joining the primitive generators of the rays of  $C$  is one. The Gorenstein index is three. By Example 3.9 any Fano polygon  $P$  with singularity content  $(n, \{m \times \frac{1}{3}(1, 1)\})$  gives rise to a toric surface  $X_P$  with degree

$$(-K_{X_P})^2 = 12 - n - \frac{5m}{3}.$$

In the notation of Definition 6.1,  $m_B = d_B = 3$  and  $s_B = 0$ .

**Theorem 7.1.** *Let  $P \subset N_{\mathbb{Q}}$  be a minimal Fano polygon with residual basket  $\mathcal{B} = \{m \times \frac{1}{3}(1, 1)\}$ , for some  $m \geq 1$ . Then, up to  $GL_2(\mathbb{Z})$ -equivalence,  $P$  is one of the 65 polygons listed in Table 4.*

*Proof.* To derive the classification, we follow the proof of Theorem 6.3.

$m_P = 3$ : Techniques for classifying all Fano polygons with given maximum Gorenstein index  $m_P$  (or with given Gorenstein index  $r$ ) are described in [28]. The resulting classifications for low index are available online [10], and it is a simple process to sift these results for the minimal polygons we require. There are precisely 60 such polygons. These are the polygons in Table 4, excluding numbers (1.1), (1.2), (1.3), (1.4), and (2.6).

$m_P > 3, k = 2$ : In this case  $P$  is a triangle with vertices  $(-3, r_E), (-3 + 2r_E, r_E)$ , and  $(-3, -r_E)$ , where  $r_E = m_P$  equals 4 or 5. We require that  $2r_E \equiv 1 \pmod{3}$ , which excludes  $r_E = 4$ . This gives polygon number (1.1) in Table 4.

$m_P = r_E > 3, k = 1$ , no edge parallel to  $E, E'$  vertical,  $l = 0, j \geq 5$ : In this case  $P$  has vertices  $(b - r_E, r_E), (b, r_E), (b, r_E - jb)$ , and is contained in the rectangle  $[b - r_E, b] \times [r_E - jb, r_E]$  where  $5 \leq j < 11$ , by (6.4),  $3 < r_E \leq 9$ , by (6.5), and  $\frac{2r_E}{j} \leq b \leq \frac{r_E}{2}$  by (6.1) and (6.2). This gives three minimal polygons: numbers (1.2), (1.3), and (2.6) in Table 4.

$m_P = r_E > 3, k = 1$ , no edge parallel to  $E, E'$  vertical,  $l > 0, j \geq 5$ : In this case  $P$  has vertices  $(b - r_E, r_E), (b, r_E), (b, r_E - jb - 1)$ , and is contained in the rectangle  $[b - r_E, b] \times [r_E - jb - 1, r_E]$  where  $5 \leq j < 11$ , by (6.10),  $3 < r_E \leq 10$ , by (6.11), and  $\frac{2r_E}{j} \leq b < \frac{r_E}{2}$ , by (6.1) and (6.6). This gives polygon number (1.4) in Table 4.  $\square$

TABLE 4. The 65 minimal Fano polygons  $P \subset N_{\mathbb{Q}}$  with singularity content  $(n, \{m \times \frac{1}{3}(1, 1)\})$ ,  $m \geq 1$ , vertices  $\mathcal{V}(P)$ , and maximum Gorenstein index  $m_P$ , up to the action of  $GL_2(\mathbb{Z})$ . The degrees  $(-K_{X_P})^2 = 12 - n - \frac{5m}{3}$  of the corresponding toric varieties are also given. The polygons are partitioned into 26 mutation-equivalence classes; the invariants  $n$  and  $m$  completely determine the mutation-equivalence class except when  $n = 6, m = 2, (-K_{X_P})^2 = \frac{8}{3}$ , and when  $n = 5, m = 1, (-K_{X_P})^2 = \frac{16}{3}$ . See Theorems 7.1 and 7.4. Those  $P$  marked with  $\star$  are chosen as the representative polygon in the mutation-equivalence class, and are depicted in Figure 2 (page 27).

#	$\mathcal{V}(P)$	$m_P$	$n$	$m$	$(-K_{X_P})^2$
$\star$ 1.1	$(7, 5), (-3, 5), (-3, -5)$	5	10	1	$\frac{1}{3}$
1.2	$(2, 5), (-3, 5), (-3, 4), (2, -11)$	5	10	1	$\frac{1}{3}$
1.3	$(2, 7), (-5, 7), (2, -7)$	7	10	1	$\frac{1}{3}$
1.4	$(3, 8), (-5, 8), (3, -8)$	8	10	1	$\frac{1}{3}$
2.1	$(11, 2), (-1, 2), (-5, -2)$	3	8	2	$\frac{2}{3}$
2.2	$(5, 2), (-1, 2), (-2, 1), (-2, -5)$	3	8	2	$\frac{2}{3}$
2.3	$(5, 2), (-1, 2), (-5, -2), (1, -2)$	3	8	2	$\frac{2}{3}$
$\star$ 2.4	$(3, 2), (-1, 2), (-5, -2), (3, -2)$	3	8	2	$\frac{2}{3}$
2.5	$(1, 2), (-1, 2), (-5, -2), (5, -2)$	3	8	2	$\frac{2}{3}$
2.6	$(2, 5), (-3, 5), (-3, 4), (1, -4), (2, -5)$	5	8	2	$\frac{2}{3}$
3.1	$(7, 2), (-1, 2), (-2, 1), (-2, -1), (-1, -2)$	3	6	3	1
$\star$ 3.2	$(3, 1), (3, 2), (-1, 2), (-2, 1), (-2, -3), (-1, -3)$	3	6	3	1
3.3	$(2, 1), (1, 2), (-1, 2), (-2, 1), (-2, -1), (2, -5)$	3	6	3	1
4.1	$(7, 2), (-1, 2), (-5, -2)$	3	9	1	$\frac{4}{3}$
$\star$ 4.2	$(3, 2), (-1, 2), (-2, 1), (-2, -3)$	3	9	1	$\frac{4}{3}$
4.3	$(1, 2), (-1, 2), (-2, 1), (1, -5)$	3	9	1	$\frac{4}{3}$
4.4	$(1, 2), (-1, 2), (-5, -2), (1, -2)$	3	9	1	$\frac{4}{3}$
5.1	$(5, 2), (-1, 2), (-2, 1), (-1, -1), (1, -2)$	3	4	4	$\frac{4}{3}$
5.2	$(3, 1), (3, 2), (-1, 2), (-2, 1), (-2, -1), (-1, -2), (1, -1)$	3	4	4	$\frac{4}{3}$
$\star$ 5.3	$(2, 1), (1, 2), (-1, 2), (-2, 1), (-2, -1), (-1, -2), (1, -2), (2, -1)$	3	4	4	$\frac{4}{3}$
6.1	$(1, 0), (2, 3), (-1, 3), (-1, -3)$	3	7	2	$\frac{5}{3}$
6.2	$(5, 2), (-1, 2), (-2, 1), (1, -2)$	3	7	2	$\frac{5}{3}$
6.3	$(5, 2), (-1, 2), (-5, -2), (-1, -1)$	3	7	2	$\frac{5}{3}$

Continued on next page.

Continued from previous page.

#	$\mathcal{V}(P)$	$m_P$	$n$	$m$	$(-K_{X_P})^2$
*6.4	$(2, 1), (1, 2), (-1, 2), (-2, 1), (-2, -3)$	3	7	2	$\frac{5}{3}$
6.5	$(1, 2), (-1, 2), (-2, 1), (-1, -1), (1, -3)$	3	7	2	$\frac{5}{3}$
*7.1	$(2, 1), (1, 2), (-1, 2), (-2, 1), (-2, -1), (-1, -2), (1, -1)$	3	2	5	$\frac{5}{3}$
8.1	$(1, 2), (-1, 2), (-2, 1), (-1, -1), (1, -2)$	3	5	3	2
*8.2	$(2, 1), (1, 2), (-1, 2), (-2, 1), (-2, -1), (-1, -2)$	3	5	3	2
*9.1	$(1, 0), (-1, 3), (-2, 3), (-1, 0), (1, -3), (2, -3)$	3	0	6	2
10.1	$(7, 1), (0, 1), (-4, -1)$	3	8	1	$\frac{7}{3}$
10.2	$(5, 2), (-1, 2), (-1, 0), (1, -2)$	3	8	1	$\frac{7}{3}$
10.3	$(1, 1), (0, 1), (-4, -1), (2, -1)$	3	8	1	$\frac{7}{3}$
10.4	$(2, 1), (0, 1), (-4, -1), (1, -1)$	3	8	1	$\frac{7}{3}$
*10.5	$(3, 1), (0, 1), (-4, -1), (0, -1)$	3	8	1	$\frac{7}{3}$
10.6	$(1, 2), (-1, 2), (-2, 1), (1, -2)$	3	8	1	$\frac{7}{3}$
*11.1	$(1, 0), (-1, 3), (-2, 3), (-1, 0), (1, -3)$	3	3	4	$\frac{7}{3}$
*12.1	$(6, 1), (0, 1), (-3, -1)$	3	6	2	$\frac{8}{3}$
12.2	$(2, 1), (0, 1), (-3, -1), (1, -1)$	3	6	2	$\frac{8}{3}$
12.3	$(3, 1), (0, 1), (-3, -1), (0, -1)$	3	6	2	$\frac{8}{3}$
12.4	$(1, 1), (0, 1), (-3, -1), (2, -1)$	3	6	2	$\frac{8}{3}$
*13.1	$(1, 0), (-1, 3), (-2, 3), (1, -3)$	3	6	2	$\frac{8}{3}$
*14.1	$(-1, 3), (-2, 3), (-1, 0), (1, -3), (1, -1)$	3	4	3	3
15.1	$(5, 1), (0, 1), (-3, -1)$	3	7	1	$\frac{10}{3}$
15.2	$(1, 1), (0, 1), (-3, -1), (1, -1)$	3	7	1	$\frac{10}{3}$
*15.3	$(2, 1), (0, 1), (-3, -1), (0, -1)$	3	7	1	$\frac{10}{3}$
15.4	$(5, 2), (-1, 2), (-1, 1), (0, -1), (1, -2)$	3	7	1	$\frac{10}{3}$
*16.1	$(-1, 3), (-2, 3), (-1, 0), (1, -2), (1, -1)$	3	5	2	$\frac{11}{3}$
*17.1	$(0, 1), (-1, 3), (-2, 3), (-1, 0), (1, -3), (1, -2)$	3	3	3	4
*18.1	$(4, 1), (0, 1), (-1, 0), (-1, -1)$	3	6	1	$\frac{13}{3}$
18.2	$(1, 0), (1, 1), (0, 1), (-3, -1), (0, -1)$	3	6	1	$\frac{13}{3}$
18.3	$(2, 1), (0, 1), (-1, 0), (-1, -1), (1, -1)$	3	6	1	$\frac{13}{3}$
18.4	$(5, 2), (-1, 2), (-5, -2), (-2, -1)$	3	6	1	$\frac{13}{3}$
18.5	$(5, 2), (-1, 2), (-1, 1), (1, -2)$	3	6	1	$\frac{13}{3}$
*19.1	$(0, 1), (-1, 3), (-2, 3), (-1, 0), (1, -2)$	3	4	2	$\frac{14}{3}$
*20.1	$(0, 1), (-1, 3), (-2, 3), (-1, 0), (1, -3)$	3	2	3	5
*21.1	$(4, 1), (0, 1), (-3, -1)$	3	5	1	$\frac{16}{3}$
21.2	$(2, 1), (0, 1), (-3, -1), (-1, -1)$	3	5	1	$\frac{16}{3}$
21.3	$(1, 1), (0, 1), (-3, -1), (0, -1)$	3	5	1	$\frac{16}{3}$
*22.1	$(3, 1), (0, 1), (-1, 0), (0, -1)$	3	5	1	$\frac{16}{3}$

Continued on next page.



Continued from previous page.


#	$\mathcal{V}(P)$	$m_P$	$n$	$m$	$(-K_{X_P})^2$
22.2	$(1, 0), (1, 1), (0, 1), (-3, -1), (-1, -1)$	3	5	1	$\frac{16}{3}$
★23.1	$(-1, 3), (-2, 3), (-1, 0), (1, -2)$	3	3	2	$\frac{17}{3}$
24.1	$(2, 1), (0, 1), (-1, 0), (1, -1)$	3	4	1	$\frac{19}{3}$
★24.2	$(1, 0), (1, 1), (0, 1), (-3, -1), (-2, -1)$	3	4	1	$\frac{19}{3}$
★25.1	$(1, 0), (1, 1), (0, 1), (-3, -1)$	3	3	1	$\frac{22}{3}$
★26.1	$(-1, 3), (-2, 3), (1, -2)$	3	2	1	$\frac{25}{3}$

We now use the minimal polygons from Theorem 7.1 to generate a complete list of mutation classes of Fano polygons with singularity content  $(n, \{m \times \frac{1}{3}(1, 1)\})$ ,  $m \geq 1$ . For future reference we recall some of the polygons from Table 4:

**Lemma 7.2.** *There are exactly two mutation-equivalence classes of Fano polygons with singularity content  $(6, \{2 \times \frac{1}{3}(1, 1)\})$ . The mutation-equivalence classes are generated by*

$$P_{12} = \text{conv}\{(6, 1), (0, 1), (-3, -1)\} \quad \text{and} \quad P_{13} = \text{conv}\{(1, 0), (-1, 3), (-2, 3), (1, -3)\}.$$

*Proof.* Notice that the six primitive  $T$ -cones in  $P_{12}$  are all contributed by the (cone over the) edge  $E$  with inner normal vector  $(0, -1) \in M$ . Hence  $\Gamma_{P_{12}}$  is a one-dimensional sublattice of  $M$ . The polygon  $P_{13}$  has primitive  $T$ -cones contributed by those edges with inner normal vectors  $(-1, 0)$  and  $(2, 1) \in M$ , hence  $\Gamma_{P_{13}}$  equals  $M$ . By Lemma 3.11 we conclude that  $P_{12}$  cannot be mutation-equivalent to  $P_{13}$ . Denote the polygons with numbers (12.2), (12.3), and (12.4) in Table 4 by  $P_{12.2}$ ,  $P_{12.3}$ , and  $P_{12.4}$  respectively. These three polygons are mutation-equivalent to  $P_{12}$ , corresponding to number (12.1) in Table 4, via:



For each mutation the primitive inner normal vector used is  $w = (0, 1)$ . Note that the left-most polygon depicted is  $GL_2(\mathbb{Z})$ -equivalent to  $P_{12}$ .  $\square$


**Lemma 7.3.** *There are exactly two mutation-equivalence classes of Fano polygons with singularity content  $(5, \{1 \times \frac{1}{3}(1, 1)\})$ . The mutation-equivalence classes are generated by*

$$P_{21} = \text{conv}\{(4, 1), (0, 1), (-3, -1)\} \quad \text{and} \quad P_{22} = \text{conv}\{(3, 1), (0, 1), (-1, 0), (0, -1)\}.$$

*Proof.* The primitive inner normal vectors to the edges of  $P_{21}$  which contribute primitive  $T$ -cones are  $(0, -1)$  and  $(-2, 7) \in M$ . These generate an index-two sublattice  $\Gamma_{P_{21}}$  of  $M$ . In the case of  $P_{22}$  the relevant inner normal vectors are  $(1, 1)$ ,  $(1, -1)$ , and  $(0, -1) \in M$ , and we have that  $[M : \Gamma_{P_{22}}] = 1$ . By Lemma 3.11 we conclude that  $P_{21}$  and  $P_{22}$  lie in distinct mutation-equivalence classes. We leave it to the reader to verify that the polygons with numbers (21.2), (21.3) (respectively, (22.2)) in Table 4 are mutation-equivalent to (21.1) (respectively, (22.1)), which is  $P_{21}$  (respectively,  $P_{22}$ ).  $\square$

**Theorem 7.4.** *There are 26 mutation-equivalence classes of Fano polygons with singularity content  $(n, \{m \times \frac{1}{3}(1, 1)\})$ ,  $m \geq 1$ . Representative polygons for each mutation-equivalence class are depicted in Figure 2 (page 27).*

*Proof.* The values  $n$  and  $m$  distinguishes every mutation class of Fano polygons with  $\frac{1}{3}(1, 1)$  singularities except in the cases  $n = 6, m = 2$  and  $n = 5, m = 1$ . These two exceptional cases are handled in Lemmas 7.2 and 7.3 above. We shall show that the minimal polygons in Table 4 with  $n = 9, m = 1$  are connected by mutation; the remaining cases are similar. There are four minimal polygons to consider, with numbers (4.1), (4.2), (4.3), and (4.4) in Table 4. Denote these polygons by  $P_{4.1}$ ,  $P_{4.2}$ ,  $P_{4.3}$ , and  $P_{4.4}$  respectively. Then, up to  $GL_2(\mathbb{Z})$ -equivalence, we have the sequence of mutations:



The mutations have been labelled with their corresponding primitive inner normal vector  $w$ .  $\square$


**Remark 7.5.** The choice of representative for each of the 26 mutation-equivalence classes depicted in Figure 2 is not arbitrary. Where possible we chose minimal polygons that are compatible with “Laurent inversion” [13], a technique for building a toric complete intersection model directly from the polygon. These models are then used by Corti–Heuberger in their classification [14] of del Pezzo surfaces with  $\frac{1}{3}(1, 1)$  singularities.

**Acknowledgements.** This work was started at the PRAGMATIC 2013 Research School in Algebraic Geometry and Commutative Algebra, “Topics in Higher Dimensional Algebraic Geometry”, held in Catania, Italy, during September 2013. We are very grateful to Alfio Ragusa, Francesco Russo, and Giuseppe Zappalà, the organisers of the PRAGMATIC school, for creating such a wonderful atmosphere in which to work. We thank Tom Coates and Alessio Corti for many useful conversations, and Daniel Cavey for pointing out that we had overlooked polygon (1.4) in Table 4 in an early draft of this paper. The majority of this paper was written during a visit by AK to Stockholm University, supported by BN’s Start-up Grant. AK is supported by EPSRC grant EP/I008128/1, ERC Starting Investigator Grant number 240123, and EPSRC Fellowship EP/N022513/1. BN is an affiliated researcher with Stockholm University and partially supported by the Vetenskapsrådet grant NT:2014-3991. TP is supported by an EPSRC Prize Studentship.

## REFERENCES

1. Mohammad Akhtar, Tom Coates, Alessio Corti, Liana Heuberger, Alexander M. Kasprzyk, Alessandro Oneto, Andrea Petracci, Thomas Prince, and Ketil Tveiten, *Mirror symmetry and the classification of orbifold del Pezzo surfaces*, Proc. Amer. Math. Soc. **144** (2016), no. 2, 513–527.
2. Mohammad Akhtar, Tom Coates, Sergey Galkin, and Alexander M. Kasprzyk, *Minkowski polynomials and mutations*, SIGMA Symmetry Integrability Geom. Methods Appl. **8** (2012), Paper 094, 17.
3. Mohammad Akhtar and Alexander M. Kasprzyk, *Singularity content*, [arXiv:1401.5458](https://arxiv.org/abs/1401.5458) [math.AG], 2014.
4. ———, *Mutations of fake weighted projective planes*, Proc. Edinb. Math. Soc. (2) **59** (2016), no. 2, 271–285.
5. Denis Auroux, *Mirror symmetry and T-duality in the complement of an anticanonical divisor*, J. Gökova Geom. Topol. GGT **1** (2007), 51–91.
6. G. Averkov, J. Krümpelmann, and B. Nill, *Largest integral simplices with one interior integral point: solution of Hensley’s conjecture and related results*, Adv. Math. **274** (2015), 118–166.
7. Victor V. Batyrev, *Higher dimensional toric varieties with ample anticanonical class*, Ph.D. thesis, Moscow State University, 1985, Text in Russian.
8. ———, *Toric degenerations of Fano varieties and constructing mirror manifolds*, The Fano Conference, Univ. Torino, Turin, 2004, pp. 109–122.
9. I. N. Bernšteĭn, I. M. Gel’fand, and V. A. Ponomarev, *Coxeter functors, and Gabriel’s theorem*, Uspehi Mat. Nauk **28** (1973), no. 2(170), 19–33.
10. Gavin Brown and Alexander M. Kasprzyk, *The Graded Ring Database*, online, access via <http://www.grdb.co.uk/>.
11. Tom Coates, Alessio Corti, Sergey Galkin, Vasily Golyshev, and Alexander M. Kasprzyk, *Mirror symmetry and Fano manifolds*, European Congress of Mathematics Kraków, 2–7 July, 2012, 2014, pp. 285–300.
12. Tom Coates, Alessio Corti, Sergey Galkin, and Alexander Kasprzyk, *Quantum periods for 3-dimensional Fano manifolds*, Geom. Topol. **20** (2016), no. 1, 103–256.
13. Tom Coates, Alexander M. Kasprzyk, and Thomas Prince, *Laurent inversion*, [arXiv:1505.01855](https://arxiv.org/abs/1505.01855) [math.AG], 2015.
14. Alessio Corti and Liana Heuberger, *Del Pezzo surfaces with  $\frac{1}{3}(1, 1)$  points*, Manuscripta Math. (2016), 1–46.

FIGURE 2. Representative minimal Fano polygons  $P \subset N_{\mathbb{Q}}$  with singularity content  $(n, \{m \times \frac{1}{3}(1,1)\})$ ,  $m \geq 1$ , for each of the 26 mutation-equivalence classes. The representatives correspond, up to the action of  $GL_2(\mathbb{Z})$ , with the polygons marked  $\star$  in Table 4. The corresponding toric varieties have degree  $(-K_{X_P})^2 = 12 - n - \frac{5m}{3}$ .



15. Dimitrios I. Dais, *Classification of toric log del Pezzo surfaces having Picard number 1 and index  $\leq 3$* , Results Math. **54** (2009), no. 3-4, 219–252.
16. V. V. Fock and A. B. Goncharov, *Cluster ensembles, quantization and the dilogarithm. II. The intertwiner*, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. I, Progr. Math., vol. 269, Birkhäuser Boston, Inc., Boston, MA, 2009, pp. 655–673.
17. Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529 (electronic).
18. William Fulton, *Introduction to toric varieties*, Ann. of Math. Stud., vol. 131, Princeton University Press, Princeton, NJ, 1993, The William H. Roever Lectures in Geometry.
19. Sergey Galkin and Alexandr Usnich, *Laurent phenomenon for ginzburg-landau potential*, IPMU preprint 10-0100, 2010.

20. Mark Gross, Paul Hacking, and Sean Keel, *Birational geometry of cluster algebras*, [arXiv:1309.2573 \[math.AG\]](https://arxiv.org/abs/1309.2573), 2013.
21. ———, *Mirror symmetry for log Calabi-Yau surfaces I*, Publ. Math. Inst. Hautes Études Sci. **122** (2015), 65–168.
22. Mark Gross and Bernd Siebert, *From real affine geometry to complex geometry*, Ann. of Math. (2) **174** (2011), no. 3, 1301–1428.
23. ———, *An invitation to toric degenerations*, Surveys in differential geometry. Volume XVI. Geometry of special holonomy and related topics, Surv. Differ. Geom., vol. 16, Int. Press, Somerville, MA, 2011, pp. 43–78.
24. Christian Haase and Josef Schicho, *Lattice polygons and the number  $2i + 7$* , Amer. Math. Monthly **116** (2009), no. 2, 151–165.
25. Paul Hacking and Yuri Prokhorov, *Smoothable del Pezzo surfaces with quotient singularities*, Compos. Math. **146** (2010), no. 1, 169–192.
26. Nathan Owen Ilten, *Mutations of Laurent polynomials and flat families with toric fibers*, SIGMA Symmetry Integrability Geom. Methods Appl. **8** (2012), 47–53.
27. Alexander M. Kasprzyk, *Bounds on fake weighted projective space*, Kodai Math. J. **32** (2009), 197–208.
28. Alexander M. Kasprzyk, Maximilian Kreuzer, and Benjamin Nill, *On the combinatorial classification of toric log del Pezzo surfaces*, LMS J. Comput. Math. **13** (2010), 33–46.
29. J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), no. 2, 299–338.
30. Maxim Kontsevich and Yan Soibelman, *Affine structures and non-Archimedean analytic spaces*, The unity of mathematics, Progr. Math., vol. 244, Birkhäuser Boston, Boston, MA, 2006, pp. 321–385.
31. Jeffrey C. Lagarias and Günter M. Ziegler, *Bounds for lattice polytopes containing a fixed number of interior points in a sublattice*, Canad. J. Math. **43** (1991), no. 5, 1022–1035.
32. Kurt Mahler, *Ein Übertragungsprinzip für konvexe Körper*, Časopis Pěst. Mat. Fys. **68** (1939), 93–102.
33. T. Mandel, *Classification of rank 2 cluster varieties*, [arXiv:1407.6241 \[math.AG\]](https://arxiv.org/abs/1407.6241), 2014.
34. Markus Perling, *Combinatorial aspects of exceptional sequences on (rational) surfaces*, [arXiv:1311.7349 \[math.AG\]](https://arxiv.org/abs/1311.7349), 2013.
35. Georg Alexander Pick, *Geometrisches zur zahlenlehre*, Sitzungber Lotos **19** (1899), 311–319.
36. Thomas Prince, *Smoothing toric Fano surfaces using the Gross–Siebert algorithm*, [arXiv:1504.05969 \[math.AG\]](https://arxiv.org/abs/1504.05969), 2015.
37. Stanley Rabinowitz, *A census of convex lattice polygons with at most one interior lattice point*, Ars Combin. **28** (1989), 83–96.
38. Miles Reid, *Young person’s guide to canonical singularities*, Algebraic geometry, Bowdoin, 1985 (Brunswick, Maine, 1985), Proc. Sympos. Pure Math., vol. 46, Amer. Math. Soc., Providence, RI, 1987, pp. 345–414.
39. P. R. Scott, *On convex lattice polygons*, Bull. Austral. Math. Soc. **15** (1976), no. 3, 395–399.
40. Margaret Symington, *Four dimensions from two in symplectic topology*, Topology and geometry of manifolds (Athens, GA, 2001), Proc. Sympos. Pure Math., vol. 71, Amer. Math. Soc., Providence, RI, 2003, pp. 153–208.
41. Jonathan M. Wahl, *Elliptic deformations of minimally elliptic singularities*, Math. Ann. **253** (1980), no. 3, 241–262.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF NOTTINGHAM, NOTTINGHAM, NG7 2RD, UK

*E-mail address:* a.m.kasprzyk@nottingham.ac.uk

DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, 180 QUEEN’S GATE, LONDON, SW7 2AZ, UK

*E-mail address:* t.prince12@imperial.ac.uk

FAKULTÄT FÜR MATHEMATIK, OTTO-VON-GUERICKE-UNIVERSITÄT MAGDEBURG, POSTSCHLISSFACH 4120, 39016 MAGDEBURG, GERMANY

*E-mail address:* benjamin.nill@ovgu.de