

Semi-orthogonal decompositions and discriminants

Ed Segal

joint with Alex Kite

Part I - Theorem

We study the derived category of coherent sheaves $D^b(X)$ on a toric variety X . The derived category has semi-orthogonal decompositions coming from wall-crossing to other birational models.

Theorem: these decompositions obey the Jordan-Hölder property.

Part II - Motivation and Conjecture

For a Calabi-Yau toric variety wall-crossing gives us many autoequivalences of $D^b(X)$. Physics/mirror symmetry predicts that these together form an action of the fundamental group of the *FI parameter space* - the complement of the discriminant in the dual toric variety.

Conjecture: the multiplicities in our decompositions agree with intersection multiplicities in the discriminant.

Semi-orthogonal decompositions

Definition

A *semi-orthogonal decomposition* of $D^b(X)$ is a sequence of full triangulated subcategories $\mathcal{C}_1, \dots, \mathcal{C}_r \subset D^b(X)$ such that:

- (i) together they generate $D^b(X)$, and
- (ii) there are no morphisms from \mathcal{C}_i to \mathcal{C}_j if $i > j$.

- Like a semi-direct product of groups.
- Gives some control over $D^b(X)$ in terms of the smaller pieces, e.g. K-theory and homology split.
- We write

$$D^b(X) = \langle \mathcal{C}_1, \dots, \mathcal{C}_r \rangle$$

If $\mathcal{C}_i \cong D^b(Y_i)$ I will write

$$D^b(X) = \langle Y_1, \dots, Y_r \rangle$$

Semi-orthogonal decompositions

Example

Let Y be a 3-fold and X be the blow-up of Y at a smooth point. Exceptional divisor is $E \cong \mathbb{P}^2$.

The sky-scraper sheaf \mathcal{O}_E is an *exceptional object* in $D^b(X)$:

$$\mathrm{End}_{D^b(X)}(\mathcal{O}_E) = \mathbb{C}$$

\implies The subcategory generated by \mathcal{O}_E is equivalent to $D^b(pt)$.

Have

$$D^b(X) = \langle Y, pt, pt \rangle$$

where the second and third subcategories are generated by $\mathcal{O}_E(2E)$ and $\mathcal{O}_E(E)$.

Semi-orthogonal decompositions

Theorem (Orlov)

Let X be the blow-up of Y in a smooth subvariety Z . Then

$$D^b(X) = \langle Y, Z, \dots, Z \rangle$$

where the number of copies of Z is $\text{codim}(Z) - 1$.

So semi-orthogonal decompositions appear when we do blow-ups.

What about other birational transformations?

Example

Let \mathbb{C}^* act on \mathbb{C}^4 with weights $(1, 1, 1, -1)$.
The two GIT quotients are:

$$X_+ \cong \mathcal{O}(-1)_{\mathbb{P}^2}$$

$$X_- \cong \mathbb{A}^3$$

We know

$$D^b(X_+) = \langle X_-, pt, pt \rangle$$

by blow-up formula.

- The *pt* here is really the fixed point (the origin) in \mathbb{C}^4 .

Example

Let \mathbb{C}^* act on \mathbb{C}^6 with weights $(1,1,1,1,-1,-1)$.
The two GIT quotients are:

$$X_+ \cong \mathcal{O}(-1)_{\mathbb{P}^2}$$

$$X_- \cong \mathbb{A}^3$$

We know

$$D^b(X_+) = \langle X_-, pt, pt \rangle$$

by blow-up formula.

- The *pt* here is really the fixed point (the origin) in \mathbb{C}^4 .

Example

Let \mathbb{C}^* act on \mathbb{C}^6 with weights $(1,1,1,1,-1,-1)$.
The two GIT quotients are:

$$X_+ \cong \mathcal{O}(-1)_{\mathbb{P}^3}^{\oplus 2}$$

$$X_- \cong \mathcal{O}(-1)_{\mathbb{P}^1}^{\oplus 4}$$

Still true that

$$D^b(X_+) = \langle X_-, pt, pt \rangle$$

- The pt here is really the fixed point (the origin) in \mathbb{C}^4 .
- The number of copies of pt equals the sum of the weights.

Theorem (Kawamata, Ballard–Favero–Katzarkov, Halpern-Leistner)

Let \mathbb{C}^* act on U . Assume $Z = U^{\mathbb{C}^*}$ connected and

$$\kappa = \text{weight}(\det(N_{Z/U})) \geq 0.$$

Then the two GIT quotients X_{\pm} obey

$$D^b(X_+) = \langle X_-, Z, \dots, Z \rangle$$

where κ copies of $D^b(Z)$ appear.

- Implies Orlov's blow-up formula.
- Could have $X_- = \emptyset$, e.g.

$$D^b(\mathbb{P}^n) = \langle pt, \dots, pt \rangle$$

where $\kappa = n + 1$ (Beilinson's theorem).


- If $\kappa = 0$ we have a flop and derived categories are equivalent.

Toric varieties

Let $(\mathbb{C}^*)^r$ act on a vector space V .

- There are many GIT quotients (“*phases*”). Each phase X_i is a toric variety.
- The space of characters has a wall-and-chamber structure, the *secondary fan*.
- A single wall crossing $X_i \rightsquigarrow X_j$ is a VGIT construction U/\mathbb{C}^* where $U \subset V$ is the semi-stable locus for a character on the wall.
- We can decompose $D^b(X_i)$ by wall-crossing repeatedly and applying theorem from previous slide.
- If X_i is compact then $D^b(X_i)$ decomposes into copies of $D^b(pt)$, if not there will be bigger pieces.

Toric varieties




Example

Let $(\mathbb{C}^*)^2$ act on \mathbb{C}^6 with weights:

$$\begin{pmatrix} 1 & 1 & -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & -1 \end{pmatrix}$$

Here $P = \mathbb{P}(\mathcal{O}^{\oplus 2} \oplus \mathcal{O}(-1))_{\mathbb{P}^1}$.

Toric varieties



Example


(1) \rightsquigarrow (2). Blows up the origin in \mathbb{A}^4 .

$$D^b(\mathcal{O}(-1)_{\mathbb{P}^3}) = \langle \mathbb{A}^4, pt, pt, pt \rangle$$

(2) \rightsquigarrow (4). Blows up $\mathcal{O}(-1)_{\mathbb{P}^1}$.

$$D^b(X) = \langle \mathcal{O}(-1)_{\mathbb{P}^3}, \mathcal{O}(-1)_{\mathbb{P}^1} \rangle = \langle \mathbb{A}^4, pt, pt, pt, \mathcal{O}(-1)_{\mathbb{P}^1} \rangle$$

Toric varieties



Example


Now go a different way. (1) \rightsquigarrow (3) blows up \mathbb{A}^2 .

$$D^b(\mathcal{O}(-1)_{\mathbb{P}^1}^1 \times \mathbb{A}^2) = \langle \mathbb{A}^4, \mathbb{A}^2 \rangle$$

(3) \rightsquigarrow (4) blows up \mathbb{P}^1 .

$$D^b(X) = \langle \mathcal{O}(-1)_{\mathbb{P}^1}^1 \times \mathbb{A}^2, \mathbb{P}^1, \mathbb{P}^1 \rangle = \langle \mathbb{A}^4, \mathbb{A}^2, \mathbb{P}^1, \mathbb{P}^1 \rangle$$

Toric varieties




Example

(1) \rightsquigarrow (2) \rightsquigarrow (4) gives $D^b(X) = \langle \mathbb{A}^4, pt, pt, pt, \mathcal{O}(-1)_{\mathbb{P}^1} \rangle$.

(1) \rightsquigarrow (3) \rightsquigarrow (4) gives $D^b(X) = \langle \mathbb{A}^4, \mathbb{A}^2, \mathbb{P}^1, \mathbb{P}^1 \rangle$.

But $\mathcal{O}(-1)_{\mathbb{P}^1}$ and \mathbb{P}^1 are toric varieties and their derived categories can also be decomposed.

Toric varieties



Example

(1) \rightsquigarrow (2) \rightsquigarrow (4) gives $D^b(X) = \langle \mathbb{A}^4, pt, pt, pt, \mathbb{A}^2, pt \rangle$.

(1) \rightsquigarrow (3) \rightsquigarrow (4) gives $D^b(X) = \langle \mathbb{A}^4, \mathbb{A}^2, pt, pt, pt, pt \rangle$.

But $\mathcal{O}(-1)_{\mathbb{P}^1}$ and \mathbb{P}^1 are toric varieties and their derived categories can also be decomposed.

Theorem (Kite-S.)

These semi-orthogonal decompositions of the derived categories of toric varieties satisfy the Jordan-Hölder property: the 'irreducible components' and their multiplicities are independent of choices.

- In the example we quotiented by $(\mathbb{C}^*)^2$ and the decomposition took 2 steps. For rank r it will take r steps.
- Proof not very hard.
- Jordan-Hölder property fails in general for semi-orthogonal decompositions [Kalck, Bondal, Böhning-Graf von Bothmer-Sosna].

Calabi-Yau toric varieties

Suppose \mathbb{C}^* acts on U and $Z = U^{\mathbb{C}^*}$ is connected and $\kappa = 0$. Then recall $D^b(X_+) \cong D^b(X_-)$. In fact the theory gives \mathbb{Z} -many equivalences:

$$\Phi_k : D^b(X_+) \xrightarrow{\sim} D^b(X_-)$$

Theorem (Halpern-Leistner–Shipman)

The autoequivalence $\Phi_1^{-1}\Phi_0$ is the twist around a spherical functor:

$$F : D^b(Z) \longrightarrow D^b(X_+)$$

If $D^b(Z)$ has a semi-orthogonal decomposition then F has a corresponding factorization.

Let $(\mathbb{C}^*)^r$ act on a vector space V through $SL(V)$.

- All phases are Calabi-Yau.
- All phases are derived equivalent.
- Wall-crossing gives many autoequivalences of each phase.
- Physics/mirror symmetry predicts:

$$\pi_1(\text{Fayet-Iliopoulos parameter space}) \curvearrowright D^b(X_i)$$

The FI parameter space is the base of the Hori-Vafa mirror
 \approx stringy Kähler moduli space of X_i .

FI parameter space

Take the dual toric variety X^\vee defined by the secondary fan. Observe:

Phases \longleftrightarrow toric fixed points in X^\vee .

Wall \longleftrightarrow toric rational curve $C_{i,j}$ connecting two fixed points.



The *FI parameter space* is the open set in X^\vee obtained by deleting:

- 1 The toric boundary.
- 2 The *GKZ discriminant locus*, a non-toric hypersurface

$$\Delta = \Delta_0 \cup \Delta_1 \cup \dots \cup \Delta_r \subset X^\vee$$

which may have several irreducible components.

FI parameter space



Loop from X_1 to X_2 and back again \rightsquigarrow the wall-crossing autoequivalence of $D^b(X_1)$.

It should factor according to (i) the components of Δ , (ii) their intersection multiplicities with $C_{1,2}$.

FI parameter space

Recall that the wall-crossing autoequivalence is the twist around a spherical functor $D^b(Z) \rightarrow D^b(X_1)$. Here Z is itself toric variety (probably not Calabi-Yau). So $D^b(Z)$ has a semi-orthogonal decomposition \implies the autoequivalence factors.

Fact: the 'irreducible components' $D^b(Y_i)$ that could occur in $D^b(Z)$ biject with the components of Δ .

Conjecture (Aspinwall-Plesser-Wang, Kite-S.)

The multiplicity of a component $D^b(Y_i) \subset D^b(Z)$ agrees with the intersection multiplicity of Δ_i with $C_{1,2}$.

Theorem (Kite-S.)

This is true in the rank 2 case.

THE END.