


Physics

Atoms ripple in the periodic table of shapes

By [Jacob Aron](#) 16 February 2011

Atoms in periodic table of shapes

Video: Atoms in periodic table of shapes



 **Unique flow pattern? It's an atom**
(Image: Imperial College London)

This rippling structure may look like a piece of origami, or an intricate scarf. In fact, it is geometry's answer to the atom because it can't be broken down into smaller components. Inspired by string theory, there is now a way to classify these atoms by their properties – and hunt down their higher dimensional cousins.

“We’re trying to build a periodic table of shapes,” says [Tom Coates](#), a mathematician at Imperial College London.

The shapes he is referring to aren’t just any old triangles or squares, but smooth ones that don’t have edges, more like spheres. These can be written as differential equations, allowing them to be described in terms of their “flow”, says Coates. If a shape has a unique flow

pattern that meets certain requirements then it is an atom. But if a shape's flow contains a number of these unique patterns, it is a molecule and can be broken down even further.

We have known for some time that such shapes can be broken down in this way. In the 1930s, Gino Fano discovered nine two-dimensional atomic shapes, while research in the 1980s revealed 102 shapes in three dimensions. But these discoveries came in the form of lists, not organised groups, and never progressed to higher dimensions.

String theory

Now Coates and his colleagues have built a computer program that can identify atoms in four and five dimensions as well as in three, and provide new insights into the properties of these atoms, such as the number of holes in each shape, or the extent to which they twist around themselves.

He now plans to use this to generate lists of shapes in higher dimensions, and then to group the shapes in each list according to their properties, much as atoms are grouped in the periodic table of elements.

Key to creating the program was [string theory](#), according to which the universe has 10 dimensions – the familiar three of space and one of time, plus another six that are hidden. The hidden dimensions may be tightly curled up in a flowing shape called the [Calabi-Yau manifold](#). And string theorists had already devised methods for turning some flowing, higher dimensional shapes into differential equations, which allow a shape's properties to be explored.

Robot arms

Coates's software uses the same mathematics to turn any smooth shape into differential equations. It then examine the shape's flow, looking for the unique patterns that signify an atom. There are hundreds of millions of potential shapes to sort through, but Coates expects a few thousand atoms.

The work could lead to insights in string theory, or help robots to make smoother motions. The path of an arm that moves forward or backward in three dimensions, or rotates along three different axes, can be represented as a higher dimensional shape. "Understanding higher dimensional shapes is very useful for motion planning," says Coates.

[David Berman](#), a string theorist at Queen Mary College, University of London, reckons the work could provide insights into superconductivity, which is already studied using string theory.

Topics: [Chemistry](#)

More from New Scientist

Explore the latest news, articles and features






Space

Seven wonders of the Milky Way: An astronomer's guide to the galaxy

Features

Subscriber-only



Environment

Dusting and vacuuming reduce wildfire air pollution inside homes

News

Free



Mathematics

AI is helping mathematicians build a periodic table of shapes

News

Subscriber-only



Comment

Lessons in Chemistry review: TV show delivers – just stay with it

Culture

Subscriber-only

Popular articles

Trending New Scientist articles

- Seven wonders of the Milky Way: An astronomer's guide to the galaxy**
- Mathematician warns US spies may be weakening next-gen encryption**
- High-dose vitamin D may lower your risk of going to hospital**
- Early humans lived in Ethiopian highlands 2 million years ago**
- Under 18s who drive independently develop a better sense of direction**
- Ukrainian AI attack drones may be killing without human oversight**
- Paris's bedbug problem is probably no worse than other major cities**
- It takes just 108 milliseconds for our brains to spot food**
- Energy-efficient transistor could allow smartwatches to use AI**
- Microgreens: The surprising truth about this trendy new 'superfood'**

Download the app



Find us on social media

Instagram

Facebook

X / Twitter

TikTok

LinkedIn

Subscriptions

Subscriber benefits

Gift

Student

Educational

Corporate

Support

Help

About us

Advertise

Write for us

Tools

Events

Science Jobs

CoLab

Syndication

RSS feeds

Legal and privacy

Contact us

Privacy policy

Cookie policy

Terms & conditions